

Quantum Group Theory in $\tau^{(2)}$ -model, Duality of $\tau^{(2)}$ -model and XXZ-model with Cyclic $U_q(sl_2)$ -representation for $q^n = 1$, and Chiral Potts Model

Shi-shyr Roan

Institute of Mathematics

Academia Sinica

Taipei, Taiwan

(email: maroan@gate.sinica.edu.tw)

We identify the quantum group $U_w(sl_2)$ in the L -operator of $\tau^{(2)}$ -model for a generic w as a subalgebra of $U_q(sl_2)$ with $w = q^{-2}$. In the roots of unity case, $q = q$, $w = \omega$ with $q^n = \omega^N = 1$, the eigenvalues and eigenvectors of XXZ-model with the $U_q(sl_2)$ -cyclic representation are determined by the $\tau^{(2)}$ -model with the induced $U_\omega(sl_2)$ -cyclic representation, which is decomposed as a finite sum of $\tau^{(2)}$ -models in non-superintegrable inhomogeneous N -state chiral Potts model. Through the theory of chiral Potts model, the Q -operator of XXZ-model can be identified with the related chiral Potts transfer matrices, with special features appeared in the $n = 2N$, e.g. N even, case. We also establish the duality of $\tau^{(2)}$ -models related to cyclic representations of $U_q(sl_2)$, analogous to the $\tau^{(2)}$ -duality in chiral Potts model; and identify the model dual to the XXZ model with $U_q(sl_2)$ -cyclic representation.

Abstract

2008 PACS: 05.50.+q, 02.20.Uw, 75.10Pq

2000 MSC: 17B37, 17B80

Key words: Quantum group, Duality of $\tau^{(2)}$ -model, XXZ-model with cyclic representation, Chiral Potts model

1 Introduction

The $\tau^{(2)}$ -model is the six-vertex model first appeared in the N -state chiral Potts model (CPM) in [11] as an adjacent model for the study of chiral Potts transfer matrix in the frame work of the Baxter's TQ -relation [5]. Together with a set of $\tau^{(j)}$ -models and functional relations among the various transfer matrices [10], one can determine all the eigenvalues of both matrices, hence solve the eigenvalue problem of CPM [1, 6, 7, 9, 20, 25]. One also expects the eigenvector problem of CPM should equally rely on the $\tau^{(2)}$ -eigenvectors as well, as already revealed in the relation of state-correspondence in the duality of chiral Potts model [27]. Furthermore, the recent progress made on the eigenvector problem in superintegrable N -state CPM for odd N in [28] again showed the vital role of $\tau^{(2)}$ -model about the degeneracy symmetries of $\tau^{(2)}$ -eigenspaces through its equivalent spin- $\frac{N-1}{2}$ XXZ-chain. It is known that the theory of quantum group $U_q(sl_2)$ for a generic q stemmed from the study of XXZ-model as an equivalent formulation of the Yang-Baxter (YB) relation (2.3) of L -operator [14, 17]. In the case when q is an N th root of unity for odd N , the XXZ chain with cyclic representation was identified with the chiral Potts $\tau^{(2)}$ -model in [26, 27]. The observation and arguments there can be further extended to a general setting for a quantum group $U_q(sl_2)$ with a generic q . In this work, we find a quantum subalgebra $U_w(sl_2)$ of $U_q(sl_2)$ for a generic $w (= q^{-2})$ (see (2.5) in the paper), with an associated L -operator satisfying the YB relation (2.6) of $\tau^{(2)}$ -model. The understanding of local state vectors of a statistical $\tau^{(2)}$ -model is thus reduced to the representation theory of $U_w(sl_2)$. The theory of N -state CPM is in essence the study of Q -operator associated to XXZ-chains with the cyclic representation in the root of unity case: $q = q$, $w = \omega$ a primitive n th and N th root of unity for $n (\geq 3), N (\geq 2)$ respectively with the relation

$$q^{-2} = \omega, \quad q^n = \omega^N = 1, \quad n = N \text{ odd, or } n = 2N. \quad (1.1)$$

The aim of this paper is to find an explicit relationship between XXZ-chains of $U_q(sl_2)$ -cyclic representation and non-superintegrable N -state chiral Potts model, with especial attention on even N case; and explore the duality theory connected to XXZ-models with cyclic representation. Since every cyclic representation of $U_q(sl_2)$ induces a representation of $U_w(sl_2)$, the XXZ-model with an $U_q(sl_2)$ -cyclic representation gives rise to a $\tau^{(2)}$ -model, denoted by $T^{(2)}(t)$ in (3.41), with the induced $U_w(sl_2)$ -cyclic representation. Correspondingly, the ABCD-algebra, i.e. the monodromy algebra, of the XXZ-model is generated by ABCD-algebra of $T^{(2)}$ -model and $K^{\frac{1}{2}}$. Hence one can study the XXZ-model with a cyclic representation through its induced $T^{(2)}$ -model. By the representation theory of $U_w(sl_2)$, we find that a $T^{(2)}$ -model with the chain-size L is decomposed as $(n/N)^L$ sub-models, each of which is isomorphic to some chiral-Potts $\tau^{(2)}$ -model with inhomogeneous vertical rapidities. The $K^{\frac{1}{2}}$ -operator of XXZ-model gives rise to a pairing of these sub-models with identical eigenvalues and eigenvectors, also with similar Q -operators. By this, the eigenvalues and eigenvectors of XXZ-model are obtained, and the Q -operator can be identified with the chiral Potts transfer matrix. In the duality theory of CPM in [27], $\tau^{(2)}$ -duality is the equality

of two $\tau^{(2)}$ -models by identifying the "ordered- and disordered-" state vectors through the spin-and-face-variable-expression of a $\tau^{(2)}$ -transfer matrix [8]. We would like to examine how far the $\tau^{(2)}$ -duality in CPM can be extended to XXZ-models with cyclic representation. First, we notice that the $\tau^{(2)}$ -duality of CPM for the sub-models in the decomposition of $T^{(2)}(t)$ can not be carried over to XXZ model, partly due to the fact that the duality works on transfer matrices only, not as ABCD-algebra- or $U_\omega(sl_2)$ -representations. However, we are still able to find a spin-and-face-expression of the transfer matrix of XXZ model with cyclic representation, and identify the model dual to it. It turns out the dual model of XXZ-chains with cyclic representation is an another type of $\tau^{(2)}$ -model, $T^{\dagger(2)}(t)$ in (3.44), dual to the $T^{(2)}(t)$ inherited from XXZ-model. This shows the dualities involved with XXZ-models all depend on the duality of $\tau^{(2)}$ -models, which we expect also serves the fundamental role in the duality of XXZ-model with a representation of $U_q(sl_2)$ other than cyclic representations.

This paper is organized as follows. In section 2, we provide a $U_W(sl_2)$ -quantum group formulation of $\tau^{(2)}$ -model for a generic w , as a parallel theory to the quantum group $U_q(sl_2)$ in XXZ-chains for a generic q . We then derive some basic properties about the monodromy-(ABCD)-algebras of the general $\tau^{(2)}$ -model. Afterwards, we shall concern only with the root of unity case (1.1) for the rest of this paper. Section 3 is devoted to the study of $\tau^{(2)}$ -model and XXZ-model with cyclic representation (of $U_\omega(sl_2)$ and $U_q(sl_2)$ respectively). In subsection 3.1, we first recall the cyclic $\mathbf{C}^n$ -representation of $U_q(sl_2)$, and describe its related cyclic representations of $U_\omega(sl_2)$; then present a detailed structure about the representation theory of L -operators in $\tau^{(2)}$ -model and XXZ-model with cyclic representation. In subsection 3.2, we study the structure of XXZ-model and $\tau^{(2)}$ -model induced from cyclic $\mathbf{C}^n$ -representation of $U_q(sl_2)$. Each irreducible component of $\tau^{(2)}$ -model with the induced cyclic $\mathbf{C}^n$ -representation is isomorphic to some $\tau^{(2)}$ -model associated to N -state CPM with (possible) inhomogeneous vertical rapidities. By this, we are able to find the connection about eigenvalues and eigenvectors between XXZ-model and $\tau^{(2)}$ -model in CPM. Section 4 is devoted to the discussion of duality of $\tau^{(2)}$ -models and XXZ models with cyclic representation. First in subsection 4.1, we recall the $\tau^{(2)}$ -duality in CPM; and then in subsection 4.2, we study the duality of $\tau^{(2)}$ -model with cyclic $\mathbf{C}^n$ -representation, and find the duality between $T^{(2)}$ - and $T^{\dagger(2)}$ -models. In subsection 4.3, we examine the relationship between the $\tau^{(2)}$ -dualities found in the previous two subsections. The $T^{(2)}$ - $T^{\dagger(2)}$ -duality agrees with the $\tau^{(2)}$ -duality in CPM only in the $n = N$ odd case, but differs in $n = 2N$ case. In subsection 4.4, we identify the $T^{\dagger(2)}$ -model as the dual model of XXZ model with cyclic representation using a spin-and-face-expression of the XXZ-transfer matrix. Section 5 is the discussion of the relationship between CPM and XXZ model with $U_q(sl_2)$ -cyclic representation. First, we recall some basic notions and the duality in CPM in subsection 5.1. Then in subsection 5.2, we identify the chiral Potts transfer matrix as the Q -operator of XXZ model with cyclic representation.

Notation: In this paper, we use the following standard notations. For a positive integer N greater than one, $\mathbf{C}^N$ denotes the vector space of N -cyclic vectors with the canonical base $|\sigma\rangle, \sigma \in \mathbf{Z}_N$ ($:= \mathbf{Z}/N\mathbf{Z}$). For a N th primitive root of unity ω , e.g. $\omega = e^{\frac{2\pi i}{N}}$, the Weyl operators

X, Z (with respect to ω) with the relations, $X^N = Z^N = 1$ and $XZ = \omega^{-1}ZX$, are defined by

$$X|\sigma\rangle = |\sigma + 1\rangle, \quad Z|\sigma\rangle = \omega^\sigma |\sigma\rangle \quad (\sigma \in \mathbf{Z}_N).$$

The Fourier basis $\{|\widehat{k}\rangle\}$ of $\{|\sigma\rangle\}$ is defined by

$$|\widehat{k}\rangle = \frac{1}{\sqrt{N}} \sum_{\sigma=0}^{N-1} \omega^{-k\sigma} |\sigma\rangle, \quad |\sigma\rangle = \frac{1}{\sqrt{N}} \sum_{k=0}^{N-1} \omega^{\sigma k} |\widehat{k}\rangle, \quad \sigma \in \mathbf{Z}_N, \quad (1.2)$$

with the corresponding Weyl operators, $\widehat{X}|\widehat{k}\rangle = |\widehat{k} + 1\rangle$, $\widehat{Z}|\widehat{k}\rangle = \omega^k |\widehat{k}\rangle$ satisfying $\widehat{X}\widehat{Z} = \omega^{-1}\widehat{Z}\widehat{X}$. Then the following equality holds:

$$(X, Z) = (\widehat{Z}, \widehat{X}^{-1}). \quad (1.3)$$

The Fourier bases of $\bigotimes^L \mathbf{C}^N$ are denoted by

$$|\sigma_1, \dots, \sigma_L\rangle := |\sigma_1\rangle \otimes \dots \otimes |\sigma_L\rangle, \quad \text{or} \quad |\widehat{k}_1, \dots, \widehat{k}_L\rangle := |\widehat{k}_1\rangle \otimes \dots \otimes |\widehat{k}_L\rangle, \quad (\sigma_j, k_j \in \mathbf{Z}_N).$$

2 Quantum Group Theory of $\tau^{(2)}$ -model

For a generic $q \in \mathbf{C} \setminus \{0, \pm 1\}$, we denote $w = q^{-2}$. It is well-known that the quantum group $U_q(sl_2)$ is the algebra generated by $K^{\pm \frac{1}{2}}, e^\pm$ with the relations $K^{\frac{1}{2}} K^{\mp \frac{1}{2}} = 1$ and

$$K^{\frac{1}{2}} e^\pm K^{-\frac{1}{2}} = q^{\pm 1} e^\pm, \quad [e^+, e^-] = \frac{K - K^{-1}}{q - q^{-1}}. \quad (2.1)$$

Define the L -operator with non-zero complex parameter $\rho, \nu \in \mathbf{C}$:

$$\mathbf{L}(s) \quad (= \mathbf{L}(s; \rho, \nu)) = \begin{pmatrix} \rho^{-1} \nu^{\frac{1}{2}} s K^{-\frac{1}{2}} - \nu^{-\frac{1}{2}} s^{-1} K^{\frac{1}{2}} & (q - q^{-1})e^- \\ (q - q^{-1})e^+ & \nu^{\frac{1}{2}} s K^{\frac{1}{2}} - \rho \nu^{-\frac{1}{2}} s^{-1} K^{-\frac{1}{2}} \end{pmatrix}. \quad (2.2)$$

The quantum group $U_q(sl_2)$ is characterized by the YB relation,

$$\mathbf{R}(s/s')(\mathbf{L}(s) \bigotimes_{aux} 1)(1 \bigotimes_{aux} \mathbf{L}(s')) = (1 \bigotimes_{aux} \mathbf{L}(s'))(\mathbf{L}(s) \bigotimes_{aux} 1)\mathbf{R}(s/s'), \quad (2.3)$$

with the symmetric R -matrix [14, 17]:

$$\mathbf{R}(s) = \begin{pmatrix} s^{-1}q - sq^{-1} & 0 & 0 & 0 \\ 0 & s^{-1} - s & q - q^{-1} & 0 \\ 0 & q - q^{-1} & s^{-1} - s & 0 \\ 0 & 0 & 0 & s^{-1}q - sq^{-1} \end{pmatrix}.$$

By setting $t = s^2$, the modified L -operator, $-s\nu^{\frac{1}{2}} K^{-\frac{1}{2}} \mathbf{L}(s)$ with the gauge transform $\text{dia}[1, -s\nu^{\frac{1}{2}}q]$, is expressed by

$$L(t) \quad (= L(t; \rho, \nu)) = \begin{pmatrix} 1 - t\nu\rho^{-1}K^{-1} & (1 - w)E^- \\ -t\nu(1 - w)E^+ & -t\nu + \rho K^{-1} \end{pmatrix}, \quad (2.4)$$

where $E^+ = -q^2 K^{\frac{-1}{2}} e^+$, $E^- = K^{\frac{-1}{2}} e^-$. The quantum subalgebra of $U_q(sl_2)$ generated by $K^{\pm 1}, E^{\pm}$, with the generator-relation:

$$KE^{\pm}K^{-1} = w^{\mp 1}E^{\pm}, \quad wE^+E^- - E^-E^+ = \frac{K^{-2}-1}{1-w}, \quad (2.5)$$

will be denoted by $U_w(sl_2)$. Then the L -matrix in (2.4) satisfy the YB equation,

$$R(t/t')(L(t) \underset{aux}{\otimes} 1)(1 \underset{aux}{\otimes} L(t')) = (1 \underset{aux}{\otimes} L(t'))(L(t) \underset{aux}{\otimes} 1)R(t/t') \quad (2.6)$$

with the asymmetry R -matrix

$$R(t) = \begin{pmatrix} tw-1 & 0 & 0 & 0 \\ 0 & t-1 & w-1 & 0 \\ 0 & t(w-1) & (t-1)w & 0 \\ 0 & 0 & 0 & tw-1 \end{pmatrix}.$$

Indeed, the generator-relation (2.5) of $U_w(sl_2)$ is characterized by the YB relation (2.6) of the L -operator (2.4). Hence a representation of $U_q(sl_2)$ or $U_w(sl_2)$ is equivalent to a representation of the L -operator (2.2) or (2.4) satisfying the respective YB relation. Two representations of L -operator (2.2) or (2.4) are equivalent if and only if the induced equivalent representations of $U_q(sl_2)$ or $U_w(sl_2)$ are equivalent with the same parameter ρ, ν . In this paper, we shall also make the identification of the quantum-algebra representation and L -operator representation when no confusion could arise.

For a chain of size L , we assign the L -operator (2.2) or (2.4) at the ℓ th site, and form the XXZ-monodromy matrix

$$\otimes_{\ell=1}^L \mathbf{L}_{\ell}(s) = \begin{pmatrix} \mathbf{A}(s) & \mathbf{B}(s) \\ \mathbf{C}(s) & \mathbf{D}(s) \end{pmatrix}, \quad \mathbf{L}_{\ell}(s) = \mathbf{L}(s; \rho_{\ell}, \nu) \text{ at site } \ell; \quad (2.7)$$

and $\tau^{(2)}$ -monodromy matrix

$$\otimes_{\ell=1}^L L_{\ell}(t) = \begin{pmatrix} A(t) & B(t) \\ C(t) & D(t) \end{pmatrix}, \quad L_{\ell}(t) = L(t; \rho_{\ell}, \nu) \text{ at site } \ell. \quad (2.8)$$

By the relation between L -operators (2.2) and (2.4), the monodromy entries of (2.7) and (2.8) are related by

$$\frac{A(t)}{D(t)} = (-s)^L \nu^{\frac{L}{2}} K^{\frac{-1}{2}} \frac{\mathbf{A}(s)}{\mathbf{D}(s)}, \quad \frac{B(t)}{C(t)} = (-s)^{L \mp 1} \nu^{\frac{L \mp 1}{2}} q^{\mp 1} K^{\frac{-1}{2}} \frac{\mathbf{B}(s)}{\mathbf{C}(s)}, \quad (2.9)$$

where $K^{\frac{1}{2}} := \bigotimes_{\ell=1}^L K_{\ell}^{\frac{1}{2}}$. Equivalently to say, the XXZ-(ABCD)-algebra (2.7) is generated by the $\tau^{(2)}$ -(ABCD)-algebra (2.8) and $K^{\frac{1}{2}}$, with the relation (2.9) between their algebra-generators. For an integer $r \in \mathbf{Z}$, the YB relation of the monodromy matrices yields the commuting relation of the r -twisted traces:

$$\mathbf{T}(s) = \mathbf{A}(s) + q^{-2r} \mathbf{D}(s), \quad \tau^{(2)}(t) = A(wt) + w^r D(wt) \quad (2.10)$$

for $s \in \mathbf{C}$ and $t \in \mathbf{C}$, as a one-parameter family in $\bigotimes^L U_q(sl_2)$ or $\bigotimes^L U_w(sl_2)$ respectively. Then we have the relation, $[\mathbf{T}(s), K^{\frac{1}{2}}] = [\tau^{(2)}(t), K] = 0$, which indeed follow from the commutative relations of $K^{\frac{1}{2}}, K$ and monodromy entries in (2.7) or (2.8):

$$K^{\frac{1}{2}} \frac{\mathbf{A}(s)}{\mathbf{D}(s)} K^{-\frac{1}{2}} = \frac{\mathbf{A}(s)}{\mathbf{D}(s)}, \quad K^{\frac{1}{2}} \frac{\mathbf{C}(s)}{\mathbf{B}(s)} K^{-\frac{1}{2}} = \frac{\mathbf{q}\mathbf{C}(s)}{\mathbf{q}^{-1}\mathbf{B}(s)}; \quad K \frac{A(t)}{D(t)} K^{-1} = \frac{A(t)}{D(t)}, \quad K \frac{C(t)}{B(t)} K^{-1} = \frac{w^{-1}C(t)}{wB(t)}.$$

By (2.9), the r -traces of $\tau^{(2)}$ and $\mathbf{T}$ in (2.10) are related by

$$\tau^{(2)}(t) = (-\mathbf{q}^{-1}s)^L \nu^{\frac{L}{2}} K^{-\frac{1}{2}} \mathbf{T}(\mathbf{q}^{-1}s), \quad t = s^2. \quad (2.11)$$

For given ρ_ℓ, ν , a $U_q(sl_2)$ -representation on $\mathbf{C}^d$ gives rise to a commuting family of transfer matrices $\mathbf{T}(s)$ or $\tau^{(2)}(t)$ on $\bigotimes^L \mathbf{C}^d$ with the L -operator induced from (2.2) or (2.4) respectively. In particular, when $\rho_\ell = 1, \nu = \mathbf{q}^{d-2}$, the spin- $\frac{d-1}{2}$ (highest weight) representation of $U_q(sl_2)$ on $\mathbf{C}^d = \bigoplus_{k=0}^{d-1} \mathbf{C}\mathbf{e}^k$:

$$\begin{aligned} K^{\frac{1}{2}}(\mathbf{e}^k) &= \mathbf{q}^{\frac{d-1-2k}{2}} \mathbf{e}^k, \quad e^+(\mathbf{e}^k) = [k]_{\mathbf{q}} \mathbf{e}^{k-1}, \quad e^-(\mathbf{e}^k) = [d-1-k]_{\mathbf{q}} \mathbf{e}^{k+1}, \\ K(\mathbf{e}^k) &= w^{\frac{-d+1}{2}+k} \mathbf{e}^k, \quad E^+(\mathbf{e}^k) = -w^{\frac{d-1}{2}-k} [k] \mathbf{e}^{k-1}, \quad E^-(\mathbf{e}^k) = w^{\frac{-d+1}{2}} [d-1-k] \mathbf{e}^{k+1}, \end{aligned} \quad (2.12)$$

where $[n]_{\mathbf{q}} := \frac{\mathbf{q}^n - \mathbf{q}^{-n}}{\mathbf{q} - \mathbf{q}^{-1}}, [n] := \frac{1-w^n}{1-w}$, for the XXZ chain (2.7) gives rise to the well-known homogeneous XXZ model of spin- $\frac{d-1}{2}$ (see, e.g. [16, 21, 22] and references therein). Note that in (2.12), $e^+(\mathbf{e}^0) = e^-(\mathbf{e}^{d-1}) = E^+(\mathbf{e}^0) = E^-(\mathbf{e}^{d-1}) = 0$.

Remark. In (2.7) and (2.8), the L -operator, $\mathbf{L}_\ell(s)$ and $L_\ell(t)$, are assumed with the same value of the parameter ν in (2.2) and (2.4). However, one may also consider monodromy matrix defined by $\mathbf{L}_\ell(s) = \mathbf{L}(s; \rho_\ell, \nu_\ell)$ and $L_\ell(t) = L(t; \rho_\ell, \nu_\ell)$ with ρ_ℓ, ν_ℓ 's distinct, and form the transfer matrix $\mathbf{T}(s; \{\rho_\ell, \nu_\ell\}), \tau^{(2)}(t; \{\rho_\ell, \nu_\ell\})$ as in (2.10). Since the L -operators in (2.2) and (2.4) satisfy the relations,

$$\mathbf{L}(s; \rho_\ell, \nu_\ell) = \mathbf{L}(\xi_\ell^{\frac{1}{2}} s; \rho_\ell, \nu), \quad L(t; \rho_\ell, \nu_\ell) = L(\xi_\ell t; \rho_\ell, \nu), \quad \nu_\ell = \xi_\ell \nu,$$

the transfer matrices $\mathbf{T}(s; \{\rho_\ell, \nu_\ell\}), \tau^{(2)}(t; \{\rho_\ell, \nu_\ell\})$ are reduced to those in (2.10) with the same ν by

$$\begin{aligned} \mathbf{T}(s; \{\rho_\ell, \nu_\ell\}) &= \mathbf{T}(\xi_1^{\frac{1}{2}} s, \dots, \xi_L^{\frac{1}{2}} s), \quad \mathbf{T}(s) = \mathbf{T}(s, \dots, s); \\ \tau^{(2)}(t; \{\rho_\ell, \nu_\ell\}) &= \tau^{(2)}(\xi_1 t, \dots, \xi_L t), \quad \tau^{(2)}(t) = \tau^{(2)}(t, \dots, t). \end{aligned} \quad (2.13)$$

3 XXZ-model and $\tau^{(2)}$ -model with Cyclic Representation

In the root of unity case (1.1), we let $|\widehat{\sigma}\rangle$'s or $|\widehat{k}\rangle$'s ($\sigma, k \in \mathbf{Z}_N$) of $\mathbf{C}^N$ be the Fourier basis in (1.2), and $(X, Z), (\widehat{X}, \widehat{Z})$ the Weyl operators (with respect to ω), in (1.3). Similarly, for the cyclic $\mathbf{n}$ -space $\mathbf{C}^{\mathbf{n}}$, the Fourier bases (with respect to q) will be denoted by

$$|\widehat{k}\rangle\rangle = \frac{1}{\sqrt{\mathbf{n}}} \sum_{\sigma=0}^{\mathbf{n}-1} q^{-k\sigma} |\sigma\rangle\rangle, \quad |\sigma\rangle\rangle = \frac{1}{\sqrt{\mathbf{n}}} \sum_{k=0}^{\mathbf{n}-1} q^{\sigma k} |\widehat{k}\rangle\rangle, \quad k, \sigma \in \mathbf{Z}_{\mathbf{n}}, \quad (3.1)$$

with the Weyl operators $(X', Z'), (\widehat{X}', \widehat{Z}')$ defined by

$$X'|\sigma\rangle\rangle = |\sigma+1\rangle\rangle, \quad Z'|\sigma\rangle\rangle = q^k |\sigma\rangle\rangle; \quad \widehat{X}'|\widehat{k}\rangle\rangle = |\widehat{k+1}\rangle\rangle, \quad \widehat{Z}'|\widehat{k}\rangle\rangle = q^k |\widehat{k}\rangle\rangle.$$

Then the relations $X'Z' = q^{-1}Z'X'$, $\widehat{X}'\widehat{Z}' = q^{-1}\widehat{Z}'\widehat{X}'$, and the identity $(X', Z') = (\widehat{Z}', \widehat{X}'^{-1})$ hold. First, we describe some special cyclic $\mathbf{C}^N$ -subspaces of $\mathbf{C}^{\mathbf{n}}$ for later use. Denote

$$c(\mathbf{n}) = 2^{\frac{-1-(-1)^{\mathbf{n}}}{4}} = \begin{cases} 1 & \text{if } \mathbf{n} = N \text{ odd} \\ 2^{-1/2} & \text{if } \mathbf{n} = 2N. \end{cases}$$

Lemma 3.1 *Consider the cyclic N -vectors in $\mathbf{C}^{\mathbf{n}}$,*

$$|\sigma\rangle\rangle_+ := c(\mathbf{n})| - 2\sigma\rangle\rangle, \quad |\sigma\rangle\rangle_- := c(\mathbf{n})| - 2\sigma + 1\rangle\rangle \in \mathbf{C}^{\mathbf{n}} \quad (\sigma \in \mathbf{Z}_N),$$

and the linear transformations

$$\wp_{\pm} : \mathbf{C}^{\mathbf{n}} \longrightarrow \mathbf{C}^N, \quad \wp_+(\widehat{|k\rangle}) = \widehat{|k\rangle}, \quad \wp_-(\widehat{|k\rangle}) = q^{-k}\widehat{|k\rangle} \quad (k \in \mathbf{Z}_{\mathbf{n}}). \quad (3.2)$$

Define the cyclic N -subspaces $\mathcal{C}^{\pm}$ of $\mathbf{C}^{\mathbf{n}}$,

$$\begin{aligned} \mathcal{C}^+ &= \sum_{\sigma=1}^{N-1} \mathbf{C}|\sigma\rangle\rangle_+ = \sum_{k=1}^{N-1} \mathbf{C}\widehat{|k\rangle}\rangle_+, \quad \widehat{|k\rangle}\rangle_+ := \frac{1}{2}(\widehat{|k\rangle}) + \widehat{|k+N\rangle}), \\ \mathcal{C}^- &= \sum_{\sigma=1}^{N-1} \mathbf{C}|\sigma\rangle\rangle_- = \sum_{k=1}^{N-1} \mathbf{C}\widehat{|k\rangle}\rangle_-, \quad \widehat{|k\rangle}\rangle_- := \frac{1}{2}(q^k\widehat{|k\rangle}) + q^{k+N}\widehat{|k+N\rangle}), \end{aligned} \quad (3.3)$$

where $\widehat{|k\rangle}\rangle_{\pm}$'s are basis of $\mathcal{C}^{\pm}$, related to $|\sigma\rangle\rangle_{\pm}$'s by the N -Fourier relation:

$$\widehat{|k\rangle}\rangle_{\pm} = \frac{1}{\sqrt{N}} \sum_{\sigma=0}^{N-1} \omega^{-k\sigma} |\sigma\rangle\rangle_{\pm}, \quad |\sigma\rangle\rangle_{\pm} = \frac{1}{\sqrt{N}} \sum_{k=0}^{N-1} \omega^{\sigma k} \widehat{|k\rangle}\rangle_{\pm}, \quad (\sigma \in \mathbf{Z}_N).$$

Then

(i) $\wp_+ = \wp_- \widehat{Z}'$, and $\widehat{Z}^{-1} \wp_- = \wp_+ \widehat{Z}'$.

(ii) The $\mathcal{C}^{\pm}$ -bases in (3.3) interchange under $\mathbf{C}^{\mathbf{n}}$ -Weyl operators via

$$X' : |\sigma\rangle\rangle_+ \mapsto |\sigma\rangle\rangle_-, \quad Z' : |\sigma\rangle\rangle_+ \mapsto \frac{\omega^{\sigma}}{q^{\omega^k}|\sigma+1\rangle\rangle_-}; \quad \widehat{X}' : \widehat{|k\rangle}\rangle_+ \mapsto \frac{\widehat{|k+1\rangle}\rangle_+}{q^{-1}\widehat{|k+1\rangle}\rangle_-}, \quad \widehat{Z}' : \widehat{|k\rangle}\rangle_+ \mapsto \frac{\widehat{|k\rangle}\rangle_-}{\omega^{-k}\widehat{|k\rangle}\rangle_+}, \quad (3.4)$$

by which $\mathbf{C}^N$ -Weyl operators of $\mathcal{C}^{\pm}$ are induced by

$$(X'^{-2}, Z'), \quad (\widehat{X}', \widehat{Z}'^{-2}) \quad \text{for } \mathcal{C}^+; \quad (X'^{-2}, q^{-1}Z'), \quad (q\widehat{X}', \widehat{Z}'^{-2}) \quad \text{for } \mathcal{C}^-. \quad (3.5)$$

Furthermore, the $\mathbf{C}^{\mathbf{n}}$ -operator X' identifies the N -spaces $\mathcal{C}^{\pm}$ by

$$X' (= \widehat{Z}') : \mathcal{C}^+ \simeq \mathcal{C}^-, \quad |\sigma\rangle\rangle_+ \mapsto |\sigma\rangle\rangle_-, \quad \widehat{|k\rangle}\rangle_+ \mapsto \widehat{|k\rangle}\rangle_-, \quad (3.6)$$

and $\mathcal{C}^{\pm}$ are isomorphic to $\mathbf{C}^N$ under $\wp_{\pm}$ in (3.2) respectively:

$$\wp_{\pm} : \mathcal{C}^{\pm} \simeq \mathbf{C}^N, \quad |\sigma\rangle\rangle_{\pm} \mapsto |\sigma\rangle, \quad \widehat{|k\rangle}\rangle_{\pm} \mapsto \widehat{|k\rangle}. \quad (3.7)$$

The automorphism (3.6) descends to the following $\mathbf{C}^N$ -automorphisms via $\wp_{\pm}$ in (3.7):

$$\mathbf{C}^N \xrightarrow{\wp_+} \mathcal{C}^+ \xrightarrow{\widehat{Z}'} \mathcal{C}^- \xrightarrow{\wp_-} \mathbf{C}^N, \quad \widehat{|k\rangle} \mapsto \widehat{|k\rangle}; \quad \mathbf{C}^N \xrightarrow{\wp_-} \mathcal{C}^- \xrightarrow{\widehat{Z}'} \mathcal{C}^+ \xrightarrow{\wp_+} \mathbf{C}^N, \quad \widehat{|k\rangle} \mapsto \omega^{-k}\widehat{|k\rangle}. \quad (3.8)$$

(iii) When $\mathbf{n} = N$ odd, $\mathbf{C}^{\mathbf{n}} = \mathcal{C}^{\pm}$ with $(|\sigma\rangle\rangle_-, \widehat{|k\rangle}\rangle_-) = (|\sigma + \frac{N-1}{2}\rangle\rangle_+, q^k\widehat{|k\rangle}\rangle_+)$, and the projections (3.7) define isomorphisms between $\mathbf{C}^{\mathbf{n}}$ and $\mathbf{C}^N$, preserving the Fourier basis. When $\mathbf{n} = 2N$, $\mathbf{C}^{\mathbf{n}} = \mathcal{C}^+ \oplus \mathcal{C}^-$, and the kernel of (3.2) $\text{Ker}(\wp_{\mp}) = \mathcal{C}^{\pm}$. Hence $|\sigma\rangle\rangle_{\pm}, \widehat{|k\rangle}\rangle_{\pm}$'s form a basis of $\mathbf{C}^{\mathbf{n}}$. In both cases, the Weyl operators of $\mathbf{C}^{\mathbf{n}}$ and $\mathbf{C}^N$ are related by

$$\wp_{\pm} X'^{-2} = X \wp_{\pm}, \quad q^{\frac{-1\pm 1}{2}} \wp_{\pm} Z' = Z \wp_{\pm}, \quad (\Leftrightarrow \quad q^{\frac{1\mp 1}{2}} \wp_{\pm} \widehat{X}' = \widehat{X} \wp_{\pm}, \quad \wp_{\pm} \widehat{Z}'^{-2} = \widehat{Z} \wp_{\pm}). \quad (3.9)$$

□

We shall also consider another kind of cyclic N -subspaces of $\mathbf{C}^{\mathbf{n}}$.

Lemma 3.2 *Consider the cyclic N -vectors in $\mathbf{C}^{\mathbf{n}}$,*

$$|\widehat{k}\rangle_+^\dagger := c(\mathbf{n})|\widehat{-2k}\rangle, \quad |\widehat{k}\rangle_-^\dagger := c(\mathbf{n})|\widehat{-2k+1}\rangle \in \mathbf{C}^{\mathbf{n}} \quad (k \in \mathbf{Z}_N),$$

and the linear transformations

$$\wp_\pm^\dagger : \mathbf{C}^{\mathbf{n}} \longrightarrow \mathbf{C}^N, \quad \wp_+^\dagger(|\sigma\rangle) = |\sigma\rangle, \quad \wp_-^\dagger(|\sigma\rangle) = q^\sigma |\sigma\rangle \quad (\sigma \in \mathbf{Z}_N). \quad (3.10)$$

Define the cyclic N -subspaces $\mathcal{C}^{\dagger\pm}$ of $\mathbf{C}^{\mathbf{n}}$,

$$\begin{aligned} \mathcal{C}^{\dagger+} &= \sum_{\sigma=1}^{N-1} \mathbf{C}|\sigma\rangle_+^\dagger = \sum_{k=1}^{N-1} \mathbf{C}|\widehat{k}\rangle_+^\dagger, \quad |\sigma\rangle_+^\dagger := \frac{1}{2}(|\sigma\rangle + |\sigma+N\rangle), \\ \mathcal{C}^{\dagger-} &= \sum_{\sigma=1}^{N-1} \mathbf{C}|\sigma\rangle_-^\dagger = \sum_{k=1}^{N-1} \mathbf{C}|\widehat{k}\rangle_-^\dagger, \quad |\sigma\rangle_-^\dagger := \frac{1}{2}(q^{-\sigma}|\sigma\rangle + q^{-\sigma-N}|\sigma+N\rangle), \end{aligned} \quad (3.11)$$

with the relation of basis:

$$|\widehat{k}\rangle_\pm^\dagger = \frac{1}{\sqrt{N}} \sum_{\sigma=0}^{N-1} \omega^{-k\sigma} |\sigma\rangle_\pm^\dagger, \quad |\sigma\rangle_\pm^\dagger = \frac{1}{\sqrt{N}} \sum_{k=0}^{N-1} \omega^{\sigma k} |\widehat{k}\rangle_\pm^\dagger, \quad (\sigma \in \mathbf{Z}_N).$$

Then

$$(i) \quad \wp_+^\dagger = \wp_-^\dagger Z'^{-1}, \text{ and } Z\wp_-^\dagger = \wp_+^\dagger Z'^{-1}.$$

(ii) The $\mathbf{C}^{\mathbf{n}}$ -Weyl operators interchange the $\mathcal{C}^{\dagger\pm}$ -basis in (3.11) by

$$X' : |\sigma\rangle_+^\dagger \mapsto |\sigma+1\rangle_+^\dagger, \quad Z' : |\sigma\rangle_+^\dagger \mapsto \omega^{-\sigma} |\sigma\rangle_-^\dagger; \quad \widehat{X}' : |\widehat{k}\rangle_+^\dagger \mapsto |\widehat{k}\rangle_-^\dagger, \quad \widehat{Z}' : |\widehat{k}\rangle_+^\dagger \mapsto \omega^k |\widehat{k}\rangle_-^\dagger, \quad (3.12)$$

by which the $\mathbf{C}^N$ -Weyl operators of $\mathcal{C}^{\dagger\pm}$ are induced by

$$(X', Z'^{-2}), \quad (\widehat{X}'^{-2}, \widehat{Z}') \text{ for } \mathcal{C}^{\dagger+}; \quad (q^{-1}X', Z'^{-2}), \quad (\widehat{X}'^{-2}, q^{-1}\widehat{Z}') \text{ for } \mathcal{C}^{\dagger-}. \quad (3.13)$$

Furthermore, the $\mathbf{C}^{\mathbf{n}}$ -operator Z'^{-1} identifies the N -spaces $\mathcal{C}^{\dagger\pm}$:

$$Z'^{-1}(= \widehat{X}') : \mathcal{C}^{\dagger+} \simeq \mathcal{C}^{\dagger-}, \quad |\sigma\rangle_+^\dagger \mapsto |\sigma\rangle_-^\dagger, \quad |\widehat{k}\rangle_+^\dagger \mapsto |\widehat{k}\rangle_-^\dagger, \quad (3.14)$$

and $\mathcal{C}^{\dagger\pm}$ are isomorphic to $\mathbf{C}^N$ under $\wp_\pm^\dagger$ in (3.10) respectively:

$$\wp_\pm^\dagger : \mathcal{C}^{\dagger\pm} \simeq \mathbf{C}^N, \quad |\sigma\rangle_\pm^\dagger \mapsto |\sigma\rangle, \quad |\widehat{k}\rangle_\pm^\dagger \mapsto |\widehat{k}\rangle. \quad (3.15)$$

The automorphism (3.14) descends to the following $\mathbf{C}^N$ -automorphisms via $\wp_\pm^\dagger$ in (3.15):

$$\mathbf{C}^N \xrightarrow{\wp_+^\dagger} \mathcal{C}^{\dagger+} \xrightarrow{\widehat{X}'} \mathcal{C}^{\dagger-} \xrightarrow{\wp_-^\dagger} \mathbf{C}^N, \quad |\sigma\rangle \mapsto |\sigma\rangle; \quad \mathbf{C}^N \xrightarrow{\wp_-^\dagger} \mathcal{C}^{\dagger-} \xrightarrow{\widehat{X}'} \mathcal{C}^{\dagger+} \xrightarrow{\wp_+^\dagger} \mathbf{C}^N, \quad |\sigma\rangle \mapsto \omega^\sigma |\sigma\rangle. \quad (3.16)$$

(iii) When $\mathbf{n} = N$ odd, $\mathbf{C}^{\mathbf{n}} = \mathcal{C}^{\dagger\pm}$ with $(|\sigma\rangle_-^\dagger, |\widehat{k}\rangle_-^\dagger) = (q^{-\sigma}|\sigma\rangle_+^\dagger, |k + \frac{N-1}{2}\rangle_+^\dagger)$, and (3.15) defines the isomorphism between $\mathbf{C}^{\mathbf{n}}$ and $\mathbf{C}^N$, preserving the Fourier basis. When $\mathbf{n} = 2N$, $\mathbf{C}^{\mathbf{n}} = \mathcal{C}^{\dagger+} \oplus \mathcal{C}^{\dagger-}$ with $\mathcal{C}^{\dagger\pm} = \text{Ker}(\wp_\mp^\dagger)$, and $|\sigma\rangle_\pm^\dagger, |\widehat{k}\rangle_\pm^\dagger$'s form a basis of $\mathbf{C}^{\mathbf{n}}$. In both cases, the Weyl operators of $\mathbf{C}^{\mathbf{n}}$ and $\mathbf{C}^N$ are related by

$$\wp_\pm^\dagger X' = q^{\frac{1\pm 1}{2}} X \wp_\pm^\dagger, \quad \wp_\pm^\dagger Z'^{-2} = Z \wp_\pm^\dagger, \quad (\Leftrightarrow \quad \wp_\pm^\dagger \widehat{X}'^{-2} = \widehat{X} \wp_\pm^\dagger, q^{\frac{-1\pm 1}{2}} \wp_\pm^\dagger \widehat{Z}' = \widehat{Z} \wp_\pm^\dagger). \quad (3.17)$$

□

3.1 The L -operators of XXZ-model and $\tau^{(2)}$ -model with cyclic representation

Consider the cyclic-representation of $U_q(sl_2)$, i.e. the three-parameter family of $U_q(sl_2)$ -representation on the cyclic space $\mathbf{C}^n$ with parameters $q^\phi, q^{-\phi'}$ and q^ε , denoted by $s_{\varepsilon, \phi, \phi'}$, and with the following expression in terms of $\mathbf{C}^n$ -Weyl operators:

$$K^{\frac{1}{2}} = q^{\frac{\phi' - \phi}{2}} \widehat{Z}', \quad e^+ = q^\varepsilon \frac{(q^{\phi+1} \widehat{Z}'^{-1} - q^{-\phi-1} \widehat{Z}') \widehat{X}'}{q - q^{-1}}, \quad e^- = q^{-\varepsilon} \frac{(q^{\phi'+1} \widehat{Z}' - q^{-\phi'-1} \widehat{Z}'^{-1}) \widehat{X}'^{-1}}{q - q^{-1}}, \quad (3.18)$$

or equivalently, a expression using the spin-operators $(X', Z') = (\widehat{Z}', \widehat{X}'^{-1})$ (see, e.g. [12, 13] or [26, 27]). By (2.2), the $U_q(sl_2)$ -representation (3.18) is equivalent to the YB relation (2.3) of the L -operator of XXZ-model:

$$\mathcal{L}(s) = \begin{pmatrix} s\rho^{-1}\nu^{\frac{1}{2}}q^{\frac{-\phi'+\phi}{2}}\widehat{Z}'^{-1} - s^{-1}\nu^{\frac{-1}{2}}q^{\frac{\phi'-\phi}{2}}\widehat{Z}' & q^{-\varepsilon}(q^{\phi'+1}\widehat{Z}' - q^{-\phi'-1}\widehat{Z}'^{-1})\widehat{X}'^{-1} \\ q^\varepsilon(q^{\phi+1}\widehat{Z}'^{-1} - q^{-\phi-1}\widehat{Z}')\widehat{X}' & s\nu^{\frac{1}{2}}q^{\frac{\phi'-\phi}{2}}\widehat{Z}' - s^{-1}\rho\nu^{\frac{-1}{2}}q^{\frac{-\phi'+\phi}{2}}\widehat{Z}'^{-1} \end{pmatrix}. \quad (3.19)$$

The above representations are all irreducible except a special one in $n = 2N$ case which is equivalent to the spin- $\frac{n-1}{2}$ representation in (2.12). The $U_q(sl_2)$ -representation (3.18) induces the $\mathbf{C}^n$ -representation of $\widehat{U}_\omega(sl_2)$ where the generators in (2.5) are expressed by

$$K = q^{\phi' - \phi} \widehat{Z}'^2, \quad E^+ = q^{\frac{-\phi - \phi'}{2} + \varepsilon} \frac{(1 - \omega^{-\phi-1} \widehat{Z}'^{-2}) \widehat{X}'}{1 - \omega}, \quad E^- = q^{\frac{\phi + \phi'}{2} - \varepsilon} \frac{(1 - \omega^{\phi'+1} \widehat{Z}'^{-2}) \widehat{X}'^{-1}}{1 - \omega}. \quad (3.20)$$

The L -operator (2.4) with the $\mathbf{C}^n$ -representation (3.20),

$$L(t) = \begin{pmatrix} 1 - t\nu\rho^{-1}q^{\phi - \phi'} \widehat{Z}'^{-2} & q^{\frac{\phi + \phi'}{2} - \varepsilon} (1 - \omega^{\phi'+1} \widehat{Z}'^{-2}) \widehat{X}'^{-1} \\ -t\nu q^{\frac{-\phi - \phi'}{2} + \varepsilon} (1 - \omega^{-\phi-1} \widehat{Z}'^{-2}) \widehat{X}' & -t\nu + \rho q^{\phi - \phi'} \widehat{Z}'^{-2} \end{pmatrix}, \quad (3.21)$$

then satisfies the YB relation (2.6). By (3.9), representations in (3.20) descend to the following three-parameter family of cyclic $\mathbf{C}^N$ -representation of $\widehat{U}_\omega(sl_2)$ via $\wp_+$ in (3.2):

$$K = q^{\phi' - \phi} \widehat{Z}^{-1}, \quad E^+ = q^{\frac{-\phi - \phi'}{2} + \varepsilon} \frac{(1 - \omega^{-\phi-1} \widehat{Z}) \widehat{X}}{1 - \omega}, \quad E^- = q^{\frac{\phi + \phi'}{2} - \varepsilon} \frac{(1 - \omega^{\phi'+1} \widehat{Z}) \widehat{X}^{-1}}{1 - \omega}. \quad (3.22)$$

By employing the cyclic representation (3.22) of $\widehat{U}_\omega(sl_2)$ on the two-parameter family of L -operator (2.4), one obtains the following five-parameter L -operators of the cyclic $\mathbf{C}^N$ -space appeared in $\tau^{(2)}$ -model of the N -state CPM [11]:

$$L(t) = \begin{pmatrix} 1 - t \frac{c}{b'b} \widehat{Z} & (\frac{1}{b} - \omega \frac{ac}{b'b} \widehat{Z}) \widehat{X}^{-1} \\ -t(\frac{1}{b'} - \frac{a'c}{b'b} \widehat{Z}) \widehat{X} & -t \frac{1}{b'b} + \omega \frac{a'ac}{b'b} \widehat{Z} \end{pmatrix}, \quad (\widehat{Z}, \widehat{X}) = (X, Z^{-1}), \quad (3.23)$$

where as in [27], the parameters a', b', a, b, c are related to $\varepsilon, \phi, \phi', \rho, \nu$ by¹:

$$q^{\varepsilon + \frac{1}{2}} = (\frac{a'b'b^3}{a})^{\frac{1}{4}}, \quad q^{\phi+1} = (\frac{a'c}{b})^{\frac{1}{2}}, \quad q^{-\phi'} = (\frac{ac}{b'})^{\frac{1}{2}}, \quad \rho = q^{-1}(\frac{a'a}{b'b})^{\frac{1}{2}}, \quad \nu = \frac{1}{b'b} \iff \\ a = \rho\nu^{-1}q^{\frac{-\phi - \phi'}{2} - \varepsilon}, \quad \omega a'a = \rho^2\nu^{-1}, \quad b = q^{\frac{-\phi - \phi'}{2} + \varepsilon}, \quad b'b = \nu^{-1}, \quad c = \rho^{-1}q^{\phi - \phi'}. \quad (3.24)$$

¹Here the parameter (a', b', a, b, c) is equal to $(\nu^{\frac{1}{2}}a', \nu^{\frac{-1}{2}}b', \nu^{\frac{-1}{2}}a, \nu^{\frac{1}{2}}b, c)$ in formula (4.17) of [27]. The difference is due to $L(t)$ in [27] is identified with the gauge of $-s\nu^{\frac{1}{2}}K^{-\frac{1}{2}}\mathcal{L}(s)$ by $\text{dia}[1, -sq]$, instead of $\text{dia}[1, -s\nu^{\frac{1}{2}}q]$ in this paper.

Note that representations in (3.22), when changing q^ε to $\omega^{-n}q^\varepsilon$, or equivalently, (a', b', a, b, c) to $(\omega^{-n}a', \omega^n b', \omega^n a, \omega^{-n}b, c)$ for $n \in \mathbf{Z}$, are equivalent. For convenience, through a factorization of c , we shall express the parameter of L -operator in (3.23) by

$$L(t; \mathbf{p}', \mathbf{p}) := L(t), \quad \mathbf{p}' = (a', b', d'), \quad \mathbf{p} = (a, b, d), \quad c = d'd. \quad (3.25)$$

We shall also write $\mathbf{p}'_+ := \mathbf{p}'$, $\mathbf{p}_+ := \mathbf{p}$. Through $\wp_-$ in (3.2) and relations in (3.9), the $U_\omega(sl_2)$ -representation (3.20) and L -operator (3.21) descend to the cyclic $\mathbf{C}^N$ -representation (3.22) with ε replaced by $\varepsilon - 1$, hence the L -operator (3.23) with parameter $(\mathbf{p}'_-, \mathbf{p}_-)$,

$$L(t; \mathbf{p}'_-, \mathbf{p}_-), \quad \mathbf{p}'_- = (a'_-, b'_-, d'_-), \quad \mathbf{p}_- = (a_-, b_-, d_-), \quad (3.26)$$

where $(\mathbf{p}'_-, \mathbf{p}_-)$ is related to $(\mathbf{p}', \mathbf{p})$ by

$$(a'_-, b'_-, a_-, b_-, c_-) = (q^{-1}a', qb', qa, q^{-1}b, c), \quad c_- = d'_-d_-, \quad (d'_-, d_-) = (d', d). \quad (3.27)$$

The relation (3.27) is equivalent to the gauge-equivalence of L -operators:

$$\begin{aligned} L(t; \mathbf{p}'_-, \mathbf{p}_-) &= \text{dia}[1, q^{-1}]L(t; \mathbf{p}'_+ \mathbf{p}_+) \text{dia}[1, q], \quad \Leftrightarrow \\ \widehat{Z}L(t; \mathbf{p}'_+, \mathbf{p}_+)\widehat{Z}^{-1} &= \text{dia}[1, q^{-1}]L(t; \mathbf{p}'_-, \mathbf{p}_-) \text{dia}[1, q]. \end{aligned} \quad (3.28)$$

The L -operators in (3.25), (3.26) are related to the $\mathbf{C}^n$ -representation in (3.20) in the following result in [26]:

Lemma 3.3 *The N -subspaces $\mathcal{C}^\pm$ of $\mathbf{C}^n$ in (3.3) are irreducible components of the $U_\omega(sl_2)$ -representation (3.20) defined by the Weyl operators in (3.5). As $U_\omega(sl_2)$ -representations, $\mathcal{C}^\pm$ is equivalent to $L(\mathbf{p}'_\pm, \mathbf{p}_\pm; t)$ via the morphism $\wp_\pm$ in (3.7). For $n = N$ odd, $\mathbf{C}^n \simeq \mathcal{C}^\pm$ with the equivalence between $\mathcal{C}^\pm$ given by $|\widehat{k}\rangle_- = q^k |\widehat{k}\rangle_+$. For $n = 2N$, $\mathbf{C}^n = \mathcal{C}^+ \oplus \mathcal{C}^-$, which is equivalent to $L(t; \mathbf{p}'_+, \mathbf{p}_+) \oplus L(t; \mathbf{p}'_-, \mathbf{p}_-)$, as $U_\omega(sl_2)$ -representations.*

Proof. It is obvious that $\mathcal{C}^\pm$ with the Weyl operators in (3.5) are irreducible components of $U_\omega(sl_2)$ -representation $\mathbf{C}^n$ in (3.20). The rest statements follows from the relations (3.7), (3.22)-(3.26), and Lemma 3.1 (iii). $\square$

Through the substitution (3.24), we shall also use the parameter $(\mathbf{p}', \mathbf{p})$ in (3.25) to represent the L -operators $\mathcal{L}(s)$, $L(t)$ in (3.19) and (3.21):

$$\begin{aligned} \mathcal{L}(s; \mathbf{p}', \mathbf{p}) &= \left(\frac{b'^3 b^3}{\omega a' a c^2} \right)^{\frac{1}{4}} \begin{pmatrix} -s^{-1} \widehat{Z}' + s \frac{c}{b' b} \widehat{Z}'^{-1} & \frac{q}{(b' b)^{\frac{1}{2}}} \left(\frac{1}{b} \widehat{Z}' - \omega \frac{a c}{b' b} \widehat{Z}'^{-1} \right) \widehat{X}'^{-1} \\ \frac{-(b' b)^{\frac{1}{2}}}{q} \left(\frac{1}{b'} \widehat{Z}' - \frac{a' c}{b' b} \widehat{Z}'^{-1} \right) \widehat{X}' & s \frac{1}{b' b} \widehat{Z}' - s^{-1} \omega \frac{a' a c}{b' b} \widehat{Z}'^{-1} \end{pmatrix}, \\ L(t; \mathbf{p}', \mathbf{p}) &= \begin{pmatrix} 1 - t \frac{c}{b' b} \widehat{Z}'^{-2} & \left(\frac{1}{b} - \omega \frac{a c}{b' b} \widehat{Z}'^{-2} \right) \widehat{X}'^{-1} \\ -t \left(\frac{1}{b'} - \frac{a' c}{b' b} \widehat{Z}'^{-2} \right) \widehat{X}' & -t \frac{1}{b' b} + \omega \frac{a' a c}{b' b} \widehat{Z}'^{-2} \end{pmatrix}, \quad (\widehat{Z}', \widehat{X}') = (X', Z'^{-1}), \end{aligned} \quad (3.29)$$

which are related by

$$\begin{aligned} L(t; \mathbf{p}' \mathbf{p}) &= \text{dia}[1, -s(b' b)^{-\frac{1}{2}} q] \left(-s \left(\frac{\omega a' a c^2}{b^3 b^3} \right)^{\frac{1}{4}} \widehat{Z}'^{-1} \mathcal{L}(s; \mathbf{p}' \mathbf{p}) \right) \text{dia}[1, -s^{-1}(b' b)^{\frac{1}{2}} q^{-1}] \Leftrightarrow \\ \mathcal{L}(s; \mathbf{p}' \mathbf{p}) &= \text{dia}[1, -s^{-1}(b' b)^{\frac{1}{2}} q^{-1}] \left(-s^{-1} \left(\frac{\omega a' a c^2}{b^3 b^3} \right)^{-\frac{1}{4}} \widehat{Z}' L(t; \mathbf{p}' \mathbf{p}) \right) \text{dia}[1, -s(b' b)^{-\frac{1}{2}} q]. \end{aligned} \quad (3.30)$$

Corresponding to (3.28), we have

$$\begin{aligned}\mathcal{L}(s; \mathbf{p}'_-, \mathbf{p}_-) &= \widehat{Z}'^{-1} \mathcal{L}(s; \mathbf{p}'_+ \mathbf{p}_+) \widehat{Z}', \quad (\Leftrightarrow \quad \mathcal{L}(s; \mathbf{p}'_+, \mathbf{p}_+) = \widehat{Z}' \mathcal{L}(s; \mathbf{p}'_-, \mathbf{p}_-) \widehat{Z}'^{-1}); \\ L(t; \mathbf{p}'_-, \mathbf{p}_-) &= \widehat{Z}'^{-1} L(t; \mathbf{p}'_+ \mathbf{p}_+) \widehat{Z}', \quad (\Leftrightarrow \quad L(t; \mathbf{p}'_+, \mathbf{p}_+) = \widehat{Z}' L(t; \mathbf{p}'_-, \mathbf{p}_-) \widehat{Z}'^{-1}).\end{aligned}\tag{3.31}$$

Indeed, by (3.16), the second relation in (3.35) is equivalent to the gauge relations in (3.28).

There is another type of cyclic $\mathbf{C}^n$ -representations of $(2.4)_{W=\omega}$ associated to the following $U_\omega(sl_2)$ representation:

$$K = q^{\phi' - \phi} \widehat{Z}'^{-1}, \quad E^+ = q^{\frac{-\phi - \phi'}{2} + \varepsilon} \frac{(1 - \omega^{-\phi - 1} \widehat{Z}') \widehat{X}'^{-2}}{1 - \omega}, \quad E^- = q^{\frac{\phi + \phi'}{2} - \varepsilon} \frac{(1 - \omega^{\phi' + 1} \widehat{Z}') \widehat{X}'^2}{1 - \omega},\tag{3.32}$$

with the L -operator

$$L^\dagger(t; \mathbf{p}', \mathbf{p}) = \begin{pmatrix} 1 - t \frac{c}{b'b} \widehat{Z}' & (\frac{1}{b} - \omega \frac{ac}{b'b} \widehat{Z}') \widehat{X}'^2 \\ -t(\frac{1}{b'} - \frac{a'c}{b'b} \widehat{Z}') \widehat{X}'^{-2} & -t \frac{1}{b'b} + \omega \frac{a'ac}{b'b} \widehat{Z}' \end{pmatrix}, \quad (\widehat{Z}', \widehat{X}') = (X', Z'^{-1}).\tag{3.33}$$

The representation (3.32) and L -operator (3.33) are obtained by the replacement of $(\widehat{X}', \widehat{Z}'^{-2})$ in (3.20), (3.29) by $(\widehat{X}'^{-2}, \widehat{Z}')$. By (3.17) and (3.24), the representation (3.32) and L -operator (3.33) descend to the $\mathbf{C}^N$ -representation (3.22) with the parameter $(\varepsilon_\pm^\dagger, \phi_\pm^\dagger, \phi_\pm^{\dagger'})$ and $L(t; \mathbf{p}_\pm^\dagger, \mathbf{p}_\pm^\dagger)$ in (3.23), via $\wp_\pm^\dagger$ in (3.10), where the parameters are defined by

$$\begin{aligned}(\varepsilon_+^\dagger, \phi_+^\dagger, \phi_+^{\dagger'}) &= (\varepsilon, \phi, \phi'), & \mathbf{p}_+^{\dagger'} &= \mathbf{p}' = (a', b', d'), \mathbf{p}_+^\dagger = \mathbf{p} = (a, b, d); \\ (\varepsilon_-^\dagger, \phi_-^\dagger, \phi_-^{\dagger'}) &= (\varepsilon, \phi + \frac{1}{2}, \phi' - \frac{1}{2}), & \mathbf{p}_-^{\dagger'} &= (a', b', d'), \mathbf{p}_-^\dagger = (a, b, dq).\end{aligned}\tag{3.34}$$

As in (3.35), we have

$$L^\dagger(t; \mathbf{p}_-^{\dagger'}, \mathbf{p}_-^\dagger) = \widehat{X}'^{-1} L^\dagger(t; \mathbf{p}_+^{\dagger'} \mathbf{p}_+^\dagger) \widehat{X}', \quad (\Leftrightarrow \quad L^\dagger(t; \mathbf{p}_+^{\dagger'}, \mathbf{p}_+^\dagger) = \widehat{X}' L^\dagger(t; \mathbf{p}_-^{\dagger'}, \mathbf{p}_-^\dagger) \widehat{X}'^{-1}),\tag{3.35}$$

which by (3.16), is equivalent to

$$L(t; \mathbf{p}_-^{\dagger'}, \mathbf{p}_-^\dagger) = \widehat{X}^{\frac{1}{2}} L(t; \mathbf{p}_+^{\dagger'} \mathbf{p}_+^\dagger) \widehat{X}^{-\frac{1}{2}}, \quad \widehat{X} L(t; \mathbf{p}_+^{\dagger'}, \mathbf{p}_+^\dagger) \widehat{X}^{-1} = \widehat{X}^{\frac{1}{2}} L(t; \mathbf{p}_-^{\dagger'}, \mathbf{p}_-^\dagger) \widehat{X}^{-\frac{1}{2}},$$

where $\widehat{X}^{-\frac{1}{2}} := \widehat{X}'$. Note that the above relation is similar to that in (3.28), but not in a form of gauge equivalence. As in Lemma 3.3, we have the following result:

Lemma 3.4 *The N -subspaces $\mathcal{C}^{\dagger\pm}$ of $\mathbf{C}^n$ in (3.11) are irreducible components of the $U_\omega(sl_2)$ -representation (3.32). As $U_\omega(sl_2)$ -representations, $\mathcal{C}^{\dagger\pm}$ is equivalent to $L(\mathbf{p}_\pm^{\dagger'}, \mathbf{p}_\pm^\dagger; t)$ via the morphism $\wp_\pm^\dagger$ in (3.15). For $n = N$ odd, $\mathbf{C}^n \simeq \mathcal{C}^{\dagger\pm}$ with the equivalence between $\mathcal{C}^{\dagger\pm}$ given by $|\sigma\rangle_-^\dagger = q^{-\sigma} |\sigma\rangle_+^\dagger$. For $n = 2N$, $\mathbf{C}^n = \mathcal{C}^{\dagger+} \oplus \mathcal{C}^{\dagger-}$ is equivalent to $L(t; \mathbf{p}_+^{\dagger'}, \mathbf{p}_+^\dagger) \oplus L(t; \mathbf{p}_-^{\dagger'}, \mathbf{p}_-^\dagger)$ as $U_\omega(sl_2)$ -representations.*

□

The following lemma describes the equivalent representations in (3.18), (3.20) or (3.32) under the change of parameter $(\mathbf{p}', \mathbf{p})$.

Lemma 3.5 We define the l -twists of $\mathbf{p} = (\mathbf{a}, \mathbf{b}, \mathbf{d})$ by

$$\mathbf{p}(l) := (\mathbf{a}q^l, \mathbf{b}q^{-l}, \mathbf{d}), \quad \mathbf{p}[l] := (\mathbf{a}, \mathbf{b}, \mathbf{d}q^l), \quad (l \in \mathbf{Z}_n). \quad (3.36)$$

Let $\mathcal{L}(s; \mathbf{p}', \mathbf{p}), L(t; \mathbf{p}', \mathbf{p})$ and $L^\dagger(t; \mathbf{p}', \mathbf{p})$ be the L -operators in (3.29), (3.33) with the parameter $(\varepsilon, \phi, \phi', \rho, \nu)$ defined in (3.24). For $l, l' \in \mathbf{Z}_n$, let $(\varepsilon^\diamond, \phi^\diamond, \phi'^\diamond, \rho^\diamond, \nu^\diamond), (\varepsilon^\circ, \phi^\circ, \phi'^\circ, \rho^\circ, \nu^\circ)$ be the parameters in (3.24) corresponding to $(\mathbf{p}'(-l'), \mathbf{p}(l))$, $(\mathbf{p}'[l'], \mathbf{p}[l])$ respectively, related to $(\varepsilon, \phi, \phi', \rho, \nu)$ by

$$\begin{aligned} q^{\varepsilon^\diamond} &= q^{\varepsilon-l}, & q^{\phi^\diamond} &= q^{\phi+\frac{l-l'}{2}}, & q^{-\phi'^\diamond} &= q^{-\phi'+\frac{l-l'}{2}}, & \rho^\diamond &= q^{l-l'}\rho, & \nu^\diamond &= q^{l-l'}\nu; \\ q^{\varepsilon^\circ} &= q^\varepsilon, & q^{\phi^\circ} &= q^{\phi+\frac{l'+l}{2}}, & q^{-\phi'^\circ} &= q^{-\phi'+\frac{l'+l}{2}}, & \rho^\circ &= \rho, & \nu^\circ &= \nu. \end{aligned} \quad (3.37)$$

Then

(i) The $\mathbf{C}^n$ -representations, $s_{\varepsilon, \phi, \phi'}$ and $s_{\varepsilon^\diamond, \phi^\diamond, \phi'^\diamond}$ of $U_q(sl_2)$ in (3.18) (or the induced $U_\omega(sl_2)$ -representation in (3.20)), are equivalent if and only if $l - l' = 2m$ ($m \in \mathbf{Z}_n$), where $s_{\varepsilon^\diamond, \phi^\diamond, \phi'^\diamond} = (\hat{X}^m \hat{Z}^{-l}) s_{\varepsilon, \phi, \phi'} (\hat{Z}^l \hat{X}^{-m})$. In particular, the L -operators in (3.29) are equivalent if and only if $l = l' \in \mathbf{Z}_n$, where $\mathcal{L}(s; \mathbf{p}'(-l), \mathbf{p}(l)) = \hat{Z}^{-l} \mathcal{L}(s; \mathbf{p}', \mathbf{p}) \hat{Z}^l$, and $L(t; \mathbf{p}'(-l), \mathbf{p}(l)) = \hat{Z}^{-l} L(t; \mathbf{p}', \mathbf{p}) \hat{Z}^l$. Furthermore, the $\mathbf{C}^N$ -representations of $U_\omega(sl_2)$ in (3.22) for $(\varepsilon, \phi, \phi')$ and $(\varepsilon^\diamond, \phi^\diamond, \phi'^\diamond)$ are equivalent if and only if $l = 2n, l - l' = 2m \in 2\mathbf{Z}_n$, where $(3.22)_{\varepsilon^\diamond, \phi^\diamond, \phi'^\diamond} = (\hat{X}^m \hat{Z}^n) (3.22)_{\varepsilon, \phi, \phi'} (\hat{Z}^{-n} \hat{X}^{-m})$. In particular, the L -operators in (3.23) are equivalent if and only if $l = l'$ with $L(t; \mathbf{p}'(-l), \mathbf{p}(l)) = \hat{Z}^n L(t; \mathbf{p}', \mathbf{p}) \hat{Z}^{-n}$.

(ii) The representations (3.32) of $U_\omega(sl_2)$ for $(\varepsilon^\circ, \phi^\circ, \phi'^\circ, \rho^\circ, \nu^\circ)$ in (3.37) are all equivalent for $l', l \in \mathbf{Z}_n$; the same for the L -operators $L^\dagger(t; \mathbf{p}'[l'], \mathbf{p}[l])$ in (3.33) with the relation $L^\dagger(t; \mathbf{p}'[l'], \mathbf{p}[l]) = \hat{X}^{-(l'+l)} L^\dagger(t; \mathbf{p}', \mathbf{p}) \hat{X}^{(l'+l)}$. Furthermore, the equivalence of $\mathbf{C}^N$ -representations (3.22) for $(\varepsilon^\circ, \phi^\circ, \phi'^\circ)$ and L -operators $L(t; \mathbf{p}'[l'], \mathbf{p}[l])$ in (3.23) is given by the condition: $l' + l = 2m \in 2\mathbf{Z}_n$, where $L(t; \mathbf{p}'[l'], \mathbf{p}[l]) = \hat{X}^m L(t; \mathbf{p}', \mathbf{p}) \hat{X}^{-m}$.

□

Remark. (I) In Lemma 3.5 (i), the representations $s_{\varepsilon^\diamond, \phi^\diamond, \phi'^\diamond}$ for $(\mathbf{p}'(-l'), \mathbf{p}(l))$ and $(\mathbf{p}'(-l'+N), \mathbf{p}(l+N))$ are equivalent with the same $\rho^\diamond, \nu^\diamond$. The parameter $(\mathbf{p}'_-, \mathbf{p}_-)$ in (3.26) is equal to $(\mathbf{p}'(-1), \mathbf{p}(1))$ with $\mathcal{L}(s), L(t)$ -relation in (3.35). When $\mathbf{n} = N$ odd, $s_{\varepsilon^\diamond, \phi^\diamond, \phi'^\diamond}$ for $l, l' \in \mathbf{Z}_n$ are all equivalent with $m = (l - l')(\frac{n+1}{2}) \in \mathbf{Z}_n$. However, in $\mathbf{n} = 2N$ case, the requirement of the constraint $l - l' = 2m$ for equivalent representations in (3.18) is necessary. Furthermore, in $\mathbf{n} = N$ odd case, one has $L(t; \mathbf{p}'(-l), \mathbf{p}(l)) = \hat{Z}^{\frac{l(N+1)}{2}} L(t; \mathbf{p}', \mathbf{p}) \hat{Z}^{-\frac{l(N+1)}{2}}$, hence $L(t; \mathbf{p}'_\pm, \mathbf{p}_\pm)$ are equivalent. When $\mathbf{n} = 2N$, since $1 \notin 2\mathbf{Z}_n$, $L(\mathbf{p}'_\pm, \mathbf{p}_\pm; t)$ are not equivalent with non-isomorphic $U_\omega(sl_2)$ -representation (3.22). (II) The $(\mathbf{p}'^\dagger_-, \mathbf{p}^\dagger_-)$ in (3.48) is equal to $(\mathbf{p}'[0], \mathbf{p}[1])$ in Lemma 3.5 (ii). Hence $L(t; \mathbf{p}'^\dagger_-, \mathbf{p}^\dagger_-) = \hat{X}^{\frac{N+1}{2}} L(t; \mathbf{p}'^\dagger_+, \mathbf{p}^\dagger_+) \hat{X}^{-\frac{(N+1)}{2}}$ for $\mathbf{n} = N$ odd case. When $\mathbf{n} = 2N$, $L(t; \mathbf{p}'^\dagger_\pm, \mathbf{p}^\dagger_\pm)$ are not equivalent.

For convenience, we shall also use the following convention to identify the index $\pm$ in Lemma 3.3 with $\mathbf{Z}_2$:

$$\gamma = \pm := \pm 1 = (-1)^i \leftrightarrow i = \frac{1-\gamma}{2} = 0, 1 \in \mathbf{Z}_2. \quad (3.38)$$

For a positive integer L , $\{\pm\}^L$ will also be identified with $\mathbf{Z}_2^L$ via

$$\vec{i} = (i_1, \dots, i_L) \in \mathbf{Z}_2^L \leftrightarrow ((-1)^{i_1}, \dots, (-1)^{i_L}) \in \{\pm\}^L,$$

in particular, $\vec{0} = (0, \dots, 0) \leftrightarrow (+, \dots, +)$, $\vec{1} = (1, \dots, 1) \leftrightarrow (-, \dots, -)$. We shall also denote $i + 1 := \vec{i} + \vec{1} \in \mathbf{Z}_2^L$.

3.2 The $\tau^{(2)}$ -model and XXZ-model with cyclic representation

For a chain of size L , we consider the $\tau^{(2)}$ -model (2.10) defined by L -operator (3.23) with parameters $\{(\mathbf{p}'_{i_\ell}, \mathbf{p}_{i_\ell})\}_{\ell=1}^L$:

$$\bigotimes_{\ell=1}^L L(t; \mathbf{p}'_{i_\ell}, \mathbf{p}_{i_\ell}) = \begin{pmatrix} A(t) & B(t) \\ C(t) & D(t) \end{pmatrix}, \quad \tau^{(2)}(t; \{\mathbf{p}'_{i_\ell}\}, \{\mathbf{p}_{i_\ell}\}) = A(t) + \omega^r D(t), \quad (3.39)$$

satisfying the boundary condition

$$|\sigma_{L+1}\rangle = |\sigma_1 - r\rangle \quad \left(\Leftrightarrow \quad |\widehat{k_{L+1}}\rangle = \omega^{-rk_1} |\widehat{k_1}\rangle \right), \quad (3.40)$$

and the periodic-parameter condition $(\mathbf{p}'_{L+1}, \mathbf{p}_{L+1}) = (\mathbf{p}'_1, \mathbf{p}_1)$. Then $\tau^{(2)}(t; \{\mathbf{p}'_{i_\ell}\}, \{\mathbf{p}_{i_\ell}\})$ commutes with the charge operator $\widehat{Z} := \prod_\ell \widehat{Z}_\ell$, which is the same as the spin-shift operator $X := \prod_\ell X_\ell$ with the eigenvalue ω^Q for $Q \in \mathbf{Z}_N$. The XXZ-model and $\tau^{(2)}$ -model in (2.10) with the monodromy matrix (2.7), (2.8) and L -operator (3.29) will be denoted by

$$\begin{aligned} \bigotimes_{\ell=1}^L \mathcal{L}(s; \mathbf{p}', \mathbf{p}) &= \begin{pmatrix} \mathcal{A}(s) & \mathcal{B}(s) \\ \mathcal{C}(s) & \mathcal{D}(s) \end{pmatrix}, \quad \mathcal{T}(s) (= \mathcal{T}(s; \mathbf{p}', \mathbf{p})) = \mathcal{A}(s) + q^{-r'} \mathcal{D}(s); \\ \bigotimes_{\ell=1}^L L(t; \mathbf{p}', \mathbf{p}) &= \begin{pmatrix} A(t) & B(t) \\ C(t) & D(t) \end{pmatrix}, \quad T^{(2)}(t) (= T^{(2)}(t; \mathbf{p}', \mathbf{p})) = A(\omega t) + q^{-r'} D(\omega t), \end{aligned} \quad (3.41)$$

satisfying the boundary condition

$$|\sigma_{L+1}\rangle = |\sigma_1 + r'\rangle \quad \left(\Leftrightarrow \quad |\widehat{k_{L+1}}\rangle = q^{r'k_1} |\widehat{k_1}\rangle \right). \quad (3.42)$$

Then $[\mathcal{T}(s), \widehat{Z}'] = [T^{(2)}(t), \widehat{Z}'^{-2}] = 0$, where $\widehat{Z}' = \prod_\ell \widehat{Z}'_\ell (= X' = \prod_\ell X'_\ell)$. The eigenvalue of $\widehat{Z}'$ and $\widehat{Z}'^{-2}$ will be denoted by $q^{Q'}, \omega^Q$ for $Q' \in \mathbf{Z}_n, Q \in \mathbf{Z}_N$ respectively. By (2.11) and (3.30), the transfer matrices in (3.41) are related by

$$T^{(2)}(t) = (-sq^{-1})^L (\frac{\omega \mathbf{a}' \mathbf{a} \mathbf{c}^2}{\mathbf{b}'^3 \mathbf{b}^3})^{\frac{L}{4}} \widehat{Z}'^{-1} \mathcal{T}(q^{-1}s) \Leftrightarrow \mathcal{T}(s) = (-s)^L (\frac{\omega \mathbf{a}' \mathbf{a} \mathbf{c}^2}{\mathbf{b}'^3 \mathbf{b}^3})^{\frac{-L}{4}} \widehat{Z}' T^{(2)}(\omega^{-1}t). \quad (3.43)$$

Similarly, we denote the $\tau^{(2)}$ -model in (2.10) with the L -operator (3.33) and the boundary condition (3.42) by

$$\bigotimes_{\ell=1}^L L^\dagger(t; \mathbf{p}', \mathbf{p}) = \begin{pmatrix} A^\dagger(t) & B^\dagger(t) \\ C^\dagger(t) & D^\dagger(t) \end{pmatrix}, \quad T^{\dagger(2)}(t) (= T^{\dagger(2)}(t; \mathbf{p}', \mathbf{p})) = A^\dagger(\omega t) + \omega^{-r'} D^\dagger(\omega t). \quad (3.44)$$

The $\widehat{Z}'^{-2}$ -eigenvalue of $T^{\dagger(2)}(t)$ will be denoted by ω^Q .

In this subsection, we study the relation between the $\tau^{(2)}$ - and XXZ-model in (3.39), (3.41) and (3.44). For $T^{(2)}(t), \mathcal{T}(s)$ in (3.41), we assume the boundary condition in (3.42) is related to (3.40) by $r' \equiv 2r \pmod{\mathbf{n}}$. First, we derive the relation between (3.39) and XXZ-model in (3.41).

For $\vec{i} = (i_1, \dots, i_L) \in \mathbf{Z}_2^L$ with $\mathbf{Z}_2$ identified with $\pm$ in (3.38), we define the following sub-quantum space of $T^{(2)}$ -model:

$$\begin{aligned} \mathcal{C}^{\vec{i}} &= \otimes_{\ell} \mathcal{C}_{\ell} \subseteq \otimes^L \mathbf{C}^{\mathbf{n}}, & (\mathcal{C}_{\ell} &:= \mathcal{C}^{(-1)^{i_{\ell}}}), \\ \wp_{\vec{i}} &(:= \otimes_{\ell} \wp_{(-1)^{i_{\ell}}}) : \mathcal{C}^{\vec{i}} \simeq \otimes^L \mathbf{C}^N, & |\widehat{k}_1, \dots, \widehat{k}_L\rangle_{\vec{i}} &:= \otimes_{\ell} |\widehat{k}_{\ell}\rangle_{(-1)^{i_{\ell}}} \mapsto |\widehat{k}_1, \dots, \widehat{k}_L\rangle, \end{aligned} \quad (3.45)$$

where $\mathcal{C}^{\pm}, \wp_{\pm}$ are in (3.3) (3.2). Then $\mathcal{C}^{\vec{i}}$ is a sub-representation of the ABCD algebra of $T^{(2)}$ -model in (3.41). By Lemma 3.3, $\wp_{\vec{i}}$ in (3.45) induces an equivalence between the $T^{(2)}$ -monodromy matrix on $\mathcal{C}^{\vec{i}}$ and $\tau^{(2)}$ -monodromy matrix (3.39) with

$$(\mathbf{p}'_{i_{\ell}}, \mathbf{p}_{i_{\ell}}) := (\mathbf{p}'_{(-1)^{i_{\ell}}}, \mathbf{p}_{(-1)^{i_{\ell}}}), \quad (3.46)$$

where $(\mathbf{p}'_{\pm}, \mathbf{p}_{\pm})$ are defined in (3.25) (3.26). Note that the parameter in (3.22) for the $\mathbf{C}^N$ -representations of $U_{\omega}(sl_2)$ associated to $L(\mathbf{p}'_{\pm}, \mathbf{p}_{\pm}; t)$ differ only in ε and $\varepsilon - 1$, in particular with the same ν in (2.8). Through $\wp_{\vec{i}}$ in (3.45), $T^{(2)}(t)$ on $\mathcal{C}^{\vec{i}}$ is equivalent to $\tau^{(2)}(t; \{\mathbf{p}'_{i_{\ell}}\}, \{\mathbf{p}_{i_{\ell}}\})$ via

$$T^{(2)}(t)|_{\mathcal{C}^{\vec{i}}} = \wp_{\vec{i}}^{-1} \cdot \tau^{(2)}(t; \{\mathbf{p}'_{i_{\ell}}\}, \{\mathbf{p}_{i_{\ell}}\}) \cdot \wp_{\vec{i}}, \quad (3.47)$$

with the relation $\widehat{Z}'^{-2} = \wp_{\vec{i}}^{-1} \cdot \widehat{Z} \cdot \wp_{\vec{i}}$. Indeed by Lemma 3.3, the structure of $T^{(2)}$ -model is given by

Proposition 3.1 *Let $T^{(2)}(t)$ be the $\tau^{(2)}$ -model in (3.41) with the boundary condition $r' = 2r$ in (3.42), and $(\mathbf{p}'_{i_{\ell}}, \mathbf{p}_{i_{\ell}})$ the parameter defined in (3.46). Then*

(i) *When $\mathbf{n} = N$ odd, $\otimes^L \mathbf{C}^{\mathbf{n}} = \mathcal{C}^{\vec{i}}$ for $\vec{i} \in \mathbf{Z}_2^L$, as representations of ABCD algebra in (2.8), and $T^{(2)}(t) \simeq \tau^{(2)}(t; \{\mathbf{p}'_{i_{\ell}}\}, \{\mathbf{p}_{i_{\ell}}\})$ via (3.47). The equivalent relations of $\mathcal{C}^{\vec{i}}$'s are induced by the isomorphism $\mathcal{C}^{\vec{0}} \simeq \mathcal{C}^{\vec{i}}: |\widehat{k}_1, \dots, \widehat{k}_L\rangle_{\vec{0}} \mapsto |\widehat{k}_1, \dots, \widehat{k}_L\rangle_{\vec{i}}$.*

(ii) *When $\mathbf{n} = 2N$, $\otimes^L \mathbf{C}^{\mathbf{n}} = \bigoplus_{\vec{i} \in \mathbf{Z}_2^L} \mathcal{C}^{\vec{i}}$ as representations of ABCD algebra in (2.8), hence relations in (3.47) give rise to the isomorphism $T^{(2)}(t) \simeq \bigoplus_{\vec{i} \in \mathbf{Z}_2^L} \tau^{(2)}(t; \{\mathbf{p}'_{i_{\ell}}\}, \{\mathbf{p}_{i_{\ell}}\})$.*

□

We now use results obtained in Proposition 3.1 to study the XXZ-model in (3.41). By (3.43), the transfer matrix $\mathcal{T}(s)$ differs from $T^{(2)}(t)$ by a scale factor and a multiple of $\widehat{Z}'$, which interchanges $\mathcal{C}^{\vec{i}}$ and $\mathcal{C}^{i\vec{i}+1}$ by (3.6). Indeed, $\widehat{Z}'$ identifies the $T^{(2)}$ -transfer matrix on these subspaces as follows:

Lemma 3.6 $T^{(2)}(t)|_{\mathcal{C}^{\vec{i}}} = \widehat{Z}'^{-1} T^{(2)}(t)|_{\mathcal{C}^{i\vec{i}+1}} \widehat{Z}'$. *As a consequence, the $T^{(2)}$ -eigenvectors $v_{\vec{i}} \in \mathcal{C}^{\vec{i}}$ and $v_{i\vec{i}+1} \in \mathcal{C}^{i\vec{i}+1}$ with the same eigenvalue are related by $v_{i\vec{i}+1} = \widehat{Z}'(v_{\vec{i}})$ (up to a non-zero scale).*

Proof. By (3.4), $\widehat{Z}'$ induces an one-to-one correspondence between $\mathcal{C}^{\vec{i}}$ and $\mathcal{C}^{i\vec{i}+1}$, which is related to the $(\otimes^L \mathbf{C}^N)$ -automorphism $\prod_{\ell} \widehat{Z}_{\ell}^{-i_{\ell}}$ via the projection $\wp_{\vec{i}}$'s in (3.45) as follows:

$$\begin{array}{ccc} \mathcal{C}^{\vec{i}} & \xrightarrow{\widehat{Z}'} & \mathcal{C}^{i\vec{i}+1}, \quad |\widehat{k}_1, \dots, \widehat{k}_L\rangle_{\vec{i}} \mapsto \omega^{-\sum_{\ell} i_{\ell} k_{\ell}} |\widehat{k}_1, \dots, \widehat{k}_L\rangle_{i\vec{i}+1} \\ \wp_{\vec{i}} \downarrow & & \downarrow \wp_{i\vec{i}+1} \quad \downarrow \\ \otimes^L \mathbf{C}^N & \xrightarrow{\prod_{\ell} \widehat{Z}_{\ell}^{-i_{\ell}}} & \otimes^L \mathbf{C}^N, \quad |\widehat{k}_1, \dots, \widehat{k}_L\rangle \mapsto \omega^{-\sum_{\ell} i_{\ell} k_{\ell}} |\widehat{k}_1, \dots, \widehat{k}_L\rangle. \end{array}$$

By (3.28), one obtains the identification

$$\tau^{(2)}(t; \{\mathbf{p}'_{i_\ell}\}, \{\mathbf{p}_{i_\ell}\}) = \left(\prod_{\ell} \widehat{Z}_{\ell}^{i_{\ell}}\right) \tau^{(2)}(t; \{\mathbf{p}'_{i_{\ell}+1}\}, \{\mathbf{p}_{i_{\ell}+1}\}) \left(\prod_{\ell} \widehat{Z}_{\ell}^{-i_{\ell}}\right).$$

Indeed, the gauge relation in (3.28) corresponds to the action of $\widehat{Z}'$ of $\bigotimes^L \mathbf{C}^{\mathbf{n}}$, hence follows the result. $\square$

Proposition 3.2 *Let $\mathcal{T}(s)$ be the XXZ-model (3.41) with the boundary condition $r' = 2r$ in (3.42), and $(\mathbf{p}'_{i_\ell}, \mathbf{p}_{i_\ell})$ be the parameters in (3.46). Then*

(i) *When $\mathbf{n} = N$ odd, $\bigotimes^L \mathbf{C}^{\mathbf{n}} = \mathcal{C}^{\vec{i}}$ for $\vec{i} \in (\mathbf{Z}_2)^L$, as representations of XXZ-(ABCD-)algebra in (2.7). The XXZ-transfer matrix $\mathcal{T}(s)$ in (3.41) is related to $\tau^{(2)}(\omega^{-1}t; \{\mathbf{p}'_{i_\ell}\}, \{\mathbf{p}_{i_\ell}\})$ by*

$$\mathcal{T}(s) = \wp_i^{-1} \cdot (-s)^L \left(\frac{\omega \mathbf{a}' \mathbf{a} \mathbf{c}^2}{\mathbf{b}'^3 \mathbf{b}^3}\right)^{\frac{-L}{4}} \widehat{Z}^{\frac{N-1}{2}} \tau^{(2)}(\omega^{-1}t; \{\mathbf{p}'_{i_\ell}\}, \{\mathbf{p}_{i_\ell}\}) \cdot \wp_i.$$

(ii) *When $\mathbf{n} = 2N$, we define $\mathcal{C}^{[\vec{i}]} = \mathcal{C}^{\vec{i}} \oplus \mathcal{C}^{i+\vec{1}}$ for a coset $[\vec{i}] := \{\vec{i}, i+\vec{1}\} \in \mathbf{Z}_2^L / \langle \vec{1} \rangle$. Then one has the decomposition of $\bigotimes^L \mathbf{C}^{\mathbf{n}}$ as representations of XXZ-algebra (2.7):*

$$\bigotimes^L \mathbf{C}^{\mathbf{n}} = \bigoplus \{\mathcal{C}^{[\vec{i}]} \mid [\vec{i}] \in \mathbf{Z}_2^L / \langle \vec{1} \rangle\}.$$

The transfer matrices $\mathcal{T}(s)$ and $\mathsf{T}^{(2)}(t)$ on each $\mathcal{C}^{[\vec{i}]}$ are related by (3.43). Indeed, $\mathcal{T}$ -eigenvectors in $\mathcal{C}^{[\vec{i}]}$ are $v_{\vec{i}} \pm q^{-Q} v_{i+\vec{1}}$ with the eigenvalue $\pm q^Q (-s)^L \left(\frac{\omega \mathbf{a}' \mathbf{a} \mathbf{c}^2}{\mathbf{b}'^3 \mathbf{b}^3}\right)^{\frac{-L}{4}} \mathsf{T}^{(2)}(\omega^{-1}t)$ and $\mathbf{Z}_{\mathbf{n}}$ -charge $Q, Q+N$ respectively, where $v_{\vec{i}}, v_{i+\vec{1}}$ are the $\mathsf{T}^{(2)}$ -eigenvectors in Lemma 3.6 with the $\mathsf{T}^{(2)}$ -eigenvalue $\mathsf{T}^{(2)}(t)$ and $\mathbf{Z}_N$ -charge Q .

Proof. (i) follows easily from Proposition 3.1 (i) and (3.43). When $\mathbf{n} = 2N$, $\widehat{Z}'$ interchanges the factors in $\mathcal{C}^{[\vec{i}]}$. Therefore $\mathcal{C}^{[\vec{i}]}$ is a component of $\bigotimes^L \mathbf{C}^{\mathbf{n}}$ as representations of XXZ-algebra. The result about $\mathcal{T}$ -eigenvectors in $\mathcal{C}^{[\vec{i}]}$ follows from Lemma 3.6, where $\mathsf{T}^{(2)}$ -eigenvectors $v_{\vec{i}}, v_{i+\vec{1}}$ are related by $\widehat{Z}'(v_{\vec{i}}) = v_{i+\vec{1}}, \widehat{Z}'(v_{i+\vec{1}}) = \omega^{-Q} v_{\vec{i}}$. $\square$

Remark. When the pair $(\vec{i}, i+\vec{1})$ is changed to $(i+\vec{1}, \vec{i})$ in Lemma 3.6, $(v_{\vec{i}}, v_{i+\vec{1}})$ are replaced by $(v_{i+\vec{1}}, \omega^{-Q} v_{\vec{i}})$, by which the $\mathcal{T}$ -eigenvectors $v_{\vec{i}} \pm q^{-Q} v_{i+\vec{1}}$ in Proposition 3.2 is changed to $v_{i+\vec{1}} + q^Q v_{\vec{i}}$.

We now study the structure of $\mathsf{T}^{\dagger(2)}$ -model in (3.44). For $\vec{i} = (i_1, \dots, i_L) \in \mathbf{Z}_2^L$, define the parameter and the sub-quantum space of $\mathsf{T}^{\dagger(2)}$ -model:

$$\begin{aligned} (\mathbf{p}_{i_\ell}^{\dagger}, \mathbf{p}_{i_\ell}^{\dagger}) &:= (\mathbf{p}_{(-1)^{i_\ell}}^{\dagger}, \mathbf{p}_{(-1)^{i_\ell}}^{\dagger}), & \mathcal{C}^{\dagger \vec{i}} &= \otimes_{\ell} \mathcal{C}_{\ell}^{\dagger} \subseteq \bigotimes^L \mathbf{C}^{\mathbf{n}}, & (\mathcal{C}_{\ell}^{\dagger} &:= \mathcal{C}^{\dagger(-1)^{i_\ell}}), \\ \wp_i^{\dagger} &:= \otimes_{\ell} \wp_{(-1)^{i_\ell}}^{\dagger} : \mathcal{C}^{\dagger \vec{i}} \simeq \bigotimes^L \mathbf{C}^N, & |\sigma_1, \dots, \sigma_L\rangle_{\vec{i}}^{\dagger} &:= \otimes_{\ell} |\sigma_{\ell}\rangle_{(-1)^{i_\ell}}^{\dagger} \mapsto |\sigma_1, \dots, \sigma_L\rangle, \end{aligned} \quad (3.48)$$

where $(\mathbf{p}_{\pm}^{\dagger}, \mathbf{p}_{\pm}^{\dagger})$ and $\mathcal{C}^{\dagger \pm}, \wp_{\pm}^{\dagger}$ are defined in (3.48), (3.11), (3.10) respectively. Similar to Proposition 3.1 and Lemma 3.6, the relation between the $\tau^{(2)}$ -models in (3.39), (3.44) is given by

Proposition 3.3 Let $T^{\dagger(2)}(t)$ be the $\tau^{(2)}$ -model in (3.44) with the boundary conditions r' in (3.42) related to r in (3.40) by $r' \equiv -r \pmod{N}$. Then $T^{\dagger(2)}(t)|_{\mathcal{C}^{\dagger\vec{i}}} = \wp_i^{\dagger-1} \cdot \tau^{\dagger(2)}(t; \{\mathbf{p}'_{i_\ell}\}, \{\mathbf{p}_{i_\ell}^\dagger\}) \cdot \wp_i^\dagger$, by which $\widehat{X}'^{-2} = \wp_i^{\dagger-1} \cdot \widehat{X} \cdot \wp_i^\dagger$, and the following results hold:

(i) When $\mathbf{n} = N$ odd, $\bigotimes^L \mathbf{C}^{\mathbf{n}} = \mathcal{C}^{\dagger\vec{i}}$ for $\vec{i} \in \mathbf{Z}_2^L$, as representations of ABCD algebra in (2.8), and $T^{\dagger(2)}(t) \simeq \tau^{(2)}(t; \{\mathbf{p}'_{i_\ell}\}, \{\mathbf{p}_{i_\ell}^\dagger\})$, where the equivalent relations among $\mathcal{C}^{\dagger\vec{i}}$'s are induced by $\mathcal{C}^{\dagger\vec{0}} \simeq \mathcal{C}^{\dagger\vec{i}}: |\sigma_1, \dots, \sigma_L\rangle_0^\dagger \mapsto |\sigma_1, \dots, \sigma_L\rangle_i^\dagger$.

(ii) When $\mathbf{n} = 2N$, $\bigotimes^L \mathbf{C}^{\mathbf{n}} = \bigoplus_{\vec{i} \in \mathbf{Z}_2^L} \mathcal{C}^{\dagger\vec{i}}$ as representations of ABCD algebra in (2.8), hence $T^{\dagger(2)}(t) \simeq \bigoplus_{\vec{i} \in \mathbf{Z}_2^L} \tau^{(2)}(t; \{\mathbf{p}'_{i_\ell}\}, \{\mathbf{p}_{i_\ell}^\dagger\})$.

(iii) $T^{\dagger(2)}(t)|_{\mathcal{C}^{\dagger\vec{i}}} = \widehat{X}'^{-1} T^{\dagger(2)}(t)|_{\mathcal{C}^{\dagger\vec{i}+1}} \widehat{X}'$, hence the $T^{\dagger(2)}$ -eigenvectors $v_i^\dagger \in \mathcal{C}^{\dagger\vec{i}}$ and $v_{i+1}^\dagger \in \mathcal{C}^{\dagger\vec{i}+1}$ with the same eigenvalue are related by $v_{i+1}^\dagger = \widehat{X}'(v_i^\dagger)$ (up to a non-zero scale).

□

4 Duality of $\tau^{(2)}$ -models and XXZ-models with cyclic representation

In this section, we discuss the duality of $\tau^{(2)}$ -models with cyclic representation, as a generalization of the $\tau^{(2)}$ -duality in CPM [27].

4.1 $\tau^{(2)}$ -duality in chiral Potts model

In this subsection, we recall the $\tau^{(2)}$ -duality in chiral Potts model in [27]. Consider the $\tau^{(2)}$ -model $\tau^{(2)}(t; \{\mathbf{p}'_{i_\ell}\}, \{\mathbf{p}_{i_\ell}\})$ in (3.39) with the boundary condition (3.40). Then $\tau^{(2)}(t; \{\mathbf{p}'_{i_\ell}\}, \{\mathbf{p}_{i_\ell}\})$ preserves the Q -subspace $V_{r,Q}$ of $\bigotimes^L \mathbf{C}^N$ with the following (Hermitian) orthonormal bases:

$$\begin{aligned} V_{r,Q} &= \bigoplus_{k_\ell} \mathbf{C}|\widehat{k}_1, \dots, \widehat{k}_L\rangle \quad (\sum_{\ell=1}^L k_\ell \equiv Q \pmod{N}, \quad \widehat{k}_{L+1} \equiv \omega^{-rk_1} \widehat{k}_1) \\ &= \bigoplus_{n_\ell} \mathbf{C}|Q; n_1, \dots, n_L\rangle \quad (\sum_{\ell=1}^L n_\ell \equiv r \pmod{N}, \quad n_{L+1} \equiv \omega^{-Qn_1} n_1), \end{aligned} \quad (4.1)$$

where $|Q; n_1, \dots, n_L\rangle := N^{-1/2} \sum_{\sigma_1=0}^{N-1} \omega^{-Q\sigma_1} |\sigma_1, \dots, \sigma_L\rangle$ with $\sigma_\ell - \sigma_{\ell+1} = n_\ell$ (see, e.g. [3, 4, 15] in $r=0$ case, and [27, 28]). Define the dual correspondence between (r, Q) - and (r^*, Q^*) -spaces (with respective to local $\mathbf{C}^N$ -basis $\{|\sigma\rangle\}, \{|\widehat{n}\rangle\}$) ([27] (3.16)):

$$\Psi: V_{r,Q} \longrightarrow V_{r^*,Q^*}, \quad |Q; n_1, \dots, n_L\rangle \mapsto |\widehat{n}_1, \dots, \widehat{n}_L\rangle, \quad (\sum_{\ell=1}^L n_\ell \equiv r) \quad (r^*, Q^*) = (Q, r); \quad (4.2)$$

and duality of parameters $\mathbf{p} \in \mathbf{C}^3$ ([27] (3.9)²):

$$\mathbf{p} = (a, b, d) \longrightarrow \mathbf{p}^* = (a^*, b^*, d^*) := (ad, bd^{-1}, d^{-1}). \quad (4.3)$$

The following lemma is used in the study of $\tau^{(2)}$ -matrix and $\tau^{(2)}$ -duality in CPM ([8] (2.14)-(2.15), [9] (3.48), [27] (3.3)-(3.9)):

²For the discussion of the chiral Potts model, the duality of rapidities in [27] (3.9) differs here by a constant $\alpha = i^{\frac{1}{N}}$ as described in Remark of Proposition 4.1 in this paper.

Lemma 4.1 *Let*

$$L(t; \mathbf{p}', \mathbf{p}) = \begin{pmatrix} L_0^0(t; \mathbf{p}', \mathbf{p}) & L_0^1(t; \mathbf{p}', \mathbf{p}) \\ L_1^0(t; \mathbf{p}', \mathbf{p}) & L_1^1(t; \mathbf{p}', \mathbf{p}) \end{pmatrix}, \quad \mathbf{p}' = (a', b', d'), \mathbf{p} = (a, b, d),$$

be L -operator in (3.23) with $\mathbf{p}', \mathbf{p}$ in (3.25). Denote

$$L_m^{m' \sigma''}(t; \mathbf{p}', \mathbf{p}) = \langle \sigma | L_m^{m'}(t; \mathbf{p}', \mathbf{p}) | \sigma'' \rangle, \quad L_n^{n' k^{*''}}(t; \mathbf{p}^*, \mathbf{p}'^*) = \langle \widehat{k^*} | L_n^{n'}(t; \mathbf{p}^*, \mathbf{p}'^*) | \widehat{k^{*''}} \rangle,$$

where $\sigma, \sigma'', k^*, k^{*''} \in \mathbf{Z}_N$, $m, m', n, n' = 0, 1$. Define

$$\begin{aligned} E(\mathbf{p}')_{m; \sigma''} &:= \omega^{-m\sigma} F_{\mathbf{p}'}(\sigma - \sigma'', m), & E(\mathbf{p})_{\sigma''; m'} &:= \omega^{m'\sigma''} \frac{\eta_{\sigma-\sigma''}}{\eta_{m'}} F_{\mathbf{p}}(\sigma - \sigma'', m') \\ U_{\mathbf{p}, \mathbf{p}'}(a|b)_c^d &:= \sum_{m=0,1} E(\mathbf{p})_a^{d; m} E(\mathbf{p}')_{m; b}^c \end{aligned} \quad (4.4)$$

for $a, b, c, d \in \mathbf{Z}_N$, where

$$\begin{aligned} \frac{\eta}{\eta_0} &= -\omega t, & F_{\mathbf{p}}(0, 0) &= 1, & F_{\mathbf{p}}(0, 1) &= \frac{-\omega t}{b}, & F_{\mathbf{p}}(1, 0) &= \frac{d}{b}, & F_{\mathbf{p}}(1, 1) &= \frac{-\omega a d}{b}, \\ & & F_{\mathbf{p}}(\alpha, m) &= 0 & \text{if } \alpha \neq 0, 1. \end{aligned} \quad (4.5)$$

Then

- (i) $L_m^{m' \sigma''}(\omega t; \mathbf{p}', \mathbf{p}) = E(\mathbf{p}')_{m; \sigma''} E(\mathbf{p})_{\sigma''; m'}$ for $\sigma, \sigma'' \in \mathbf{Z}_N$, $m, m' = 0, 1$.
- (ii) $U_{\mathbf{p}, \mathbf{p}'}(a|b)_c^d = 0$ if $a - d$ or $b - c \neq 0, 1$, and

$$U_{\mathbf{p}, \mathbf{p}'}(a|b)_c^d = L_n^{n' k^{*''}}(\omega t; \mathbf{p}^*, \mathbf{p}'^*) \quad \text{when } (a - d, b - c) = (n, n'), n, n' = 0, 1,$$

where $a - b = k^*, d - c = k^{*''} \in \mathbf{Z}_N$.

Proof. (i) follows from (3.23) and the definition of $E(\mathbf{p}')_{m; \sigma''}$ and $E(\mathbf{p})_{\sigma''; m'}$. One also finds

$$U_{\mathbf{p}, \mathbf{p}'}(a|b)_c^d = \sum_{m=0,1} \omega^{m(d-b)} (-\omega t)^{a-d-m} F_{\mathbf{p}}(a-d, m) F_{\mathbf{p}'}(b-c, m)$$

whose non-zero values are determined by entries of

$$L(\omega t; \mathbf{p}^*, \mathbf{p}'^*) = \begin{pmatrix} 1 - \omega t \frac{1}{bb'} \widehat{Z} & (\frac{d'}{b'} - \omega \frac{ad'}{bb'}) \widehat{Z} \widehat{X}^{-1} \\ -\omega t (\frac{d}{b} - \frac{ad}{bb'} \widehat{Z}) \widehat{X} & -\omega t \frac{dd'}{bb'} + \omega \frac{aa'dd'}{bb'} \widehat{Z} \end{pmatrix}.$$

Then (ii) follows. $\square$

Using Lemma 4.1 and (4.4), one finds the product form of $\tau^{(2)}(t; \{\mathbf{p}'_{i_\ell}\}, \{\mathbf{p}_{i_\ell}\})$ ([9] (3.44a) $_{k=0, j=2}$, [8] (2.16)):

$$\tau^{(2)}(t; \{\mathbf{p}'_{i_\ell}\}, \{\mathbf{p}_{i_\ell}\})_{\{\sigma_\ell\}}^{\{\sigma''_\ell\}} = \prod_{\ell=1}^L U_{\mathbf{p}_\ell, \mathbf{p}'_{\ell+1}}(\sigma''_\ell | \sigma''_{\ell+1}), \quad (4.6)$$

where $U_{\mathbf{p}, \mathbf{p}'}(a|b)_c^d$ are defined in (4.4). Note that by the boundary condition (3.40), $E(\mathbf{p}'_{L+1})_{m; \sigma''_{L+1}} = \omega^{mr} E(\mathbf{p}'_1)_{m; \sigma''_1}$, which contributes the ω^r -factor in (3.39).

Proposition 4.1 ([27] Proposition 3.1) *Let $\mathbf{p}_\ell^*, \mathbf{p}'_\ell^*$ be the dual of $\mathbf{p}_\ell, \mathbf{p}'_\ell$ in (4.3), and Ψ be the dual correspondence between $V_{r, Q}$ and V_{r^*, Q^*} with $(r^*, Q^*) = (Q, r)$ in (4.2). Then $\tau^{(2)}(t; \{\mathbf{p}'_{i_\ell}\}, \{\mathbf{p}_{i_\ell}\})$ on $V_{r, Q}$ is equivalent to $\tau^{(2)}(t; \{\mathbf{p}_{i_\ell}^*\}, \{\mathbf{p}'_{i_\ell+1}^*\})$ on V_{r^*, Q^*} by*

$$\tau^{(2)}(t; \{\mathbf{p}'_{i_\ell}\}, \{\mathbf{p}_{i_\ell}\}) = \Psi^{-1} \tau^{(2)}(t; \{\mathbf{p}_{i_\ell}^*\}, \{\mathbf{p}'_{i_\ell+1}^*\}) \Psi. \quad (4.7)$$

Proof. By (4.6) and Lemma 4.1, one finds

$$\langle Q; n_1, \dots, n_L | \tau^{(2)}(t; \{\mathbf{p}'_{i_\ell}\}, \{\mathbf{p}_{i_\ell}\}) | Q; n'_1, \dots, n'_L \rangle = \langle \hat{n}_1, \dots, \hat{n}_L | \tau^{(2)}(t; \{\mathbf{p}^*_{i_\ell}\}, \{\mathbf{p}'^*_{i_\ell+1}\}) | \hat{n}'_1, \dots, \hat{n}'_L \rangle.$$

Then the result follows. $\square$

Remark. One may modify the duality of parameters in (4.3) by defining $\mathbf{p}^* = (\alpha \mathbf{ad}, \alpha \mathbf{bd}^{-1}, \mathbf{d}^{-1})$, where α is a scale constant. Then $U_{\mathbf{p}, \mathbf{p}'}(d|_b^c)_{\text{in}} \text{ in (4.4)} = \text{dia}[1, \alpha^{-1}] L(\omega t^*; \mathbf{p}^*, \mathbf{p}'^*) \text{dia}[1, \alpha]$ with $t^* = \alpha^2 t$, by which the duality (4.7) again holds if $\tau^{(2)}(t; \{\mathbf{p}^*_{i_\ell}\}, \{\mathbf{p}'^*_{i_\ell+1}\})$ in (4.7) is changed to $\tau^{(2)}(t^*; \{\mathbf{p}^*_{i_\ell}\}, \{\mathbf{p}'^*_{i_\ell+1}\})$.

4.2 Duality of $\mathbf{t}^{(2)}$ -models with cyclic representation

We now study the duality of $\tau^{(2)}$ -models $\mathbf{T}^{(2)}(t; \mathbf{p}', \mathbf{p})$, $\mathbf{T}^{\dagger(2)}(t; \mathbf{p}', \mathbf{p})$ in (3.41), (3.44) with the boundary condition (3.42). As in (4.1), these $\tau^{(2)}$ -models preserve the Q' -subspace $V'_{r', Q'}$ of $\bigotimes^L \mathbf{C}^{\mathbf{n}}$, which is generated by the following orthonormal bases:

$$\begin{aligned} V'_{r', Q'} &= \bigoplus_{k_\ell} \mathbf{C} | \hat{k}_1, \dots, \hat{k}_L \rangle \quad (\sum_{\ell=1}^L k_\ell \equiv Q' \pmod{\mathbf{n}}, \hat{k}_{L+1} \equiv q^{r' k_1} \hat{k}_1) \\ &= \bigoplus_{n_\ell} \mathbf{C} | Q'; n_1, \dots, n_L \rangle \quad (\sum_{\ell=1}^L n_\ell \equiv -r' \pmod{\mathbf{n}}, n_{L+1} \equiv q^{-Q' n_1} n_1), \end{aligned} \quad (4.8)$$

where $|Q'; n_1, \dots, n_L \rangle := \mathbf{n}^{-1/2} \sum_{\sigma_1=0}^{\mathbf{n}-1} q^{-Q' \sigma_1} | \sigma_1, \dots, \sigma_L \rangle$ with $\sigma_\ell - \sigma_{\ell+1} = n_\ell$. There is the dual correspondence between (r', Q') - and (r'^*, Q'^*) -spaces (with respective to the local $\mathbf{C}^{\mathbf{n}}$ -basis $\{|\sigma\rangle\}, \{|\hat{n}\rangle\}$) with $(r'^*, Q'^*) = (-Q, -r)$:

$$\Psi' : V'_{r', Q'} \longrightarrow V'_{r'^*, Q'^*}, \quad |Q'; n_1, \dots, n_L \rangle \mapsto |\hat{n}_1, \dots, \hat{n}_L \rangle, \quad (\sum_{\ell=1}^L n_\ell \equiv -r \pmod{\mathbf{n}}). \quad (4.9)$$

As the $\tau^{(2)}$ -model in Lemma 4.1, the L -operator of $\mathbf{T}^{(2)}$ - and $\mathbf{T}^{\dagger(2)}$ -model can be decomposed into a product form:

Lemma 4.2 *Let*

$$\mathbf{L}(t; \mathbf{p}', \mathbf{p}) = \begin{pmatrix} \mathbf{L}_0^0(t; \mathbf{p}', \mathbf{p}) & \mathbf{L}_0^1(t; \mathbf{p}', \mathbf{p}) \\ \mathbf{L}_1^0(t; \mathbf{p}', \mathbf{p}) & \mathbf{L}_1^1(t; \mathbf{p}', \mathbf{p}) \end{pmatrix}, \quad \mathbf{L}^\dagger(t; \mathbf{p}', \mathbf{p}) = \begin{pmatrix} \mathbf{L}_0^{\dagger 0}(t; \mathbf{p}', \mathbf{p}) & \mathbf{L}_0^{\dagger 1}(t; \mathbf{p}', \mathbf{p}) \\ \mathbf{L}_1^{\dagger 0}(t; \mathbf{p}', \mathbf{p}) & \mathbf{L}_1^{\dagger 1}(t; \mathbf{p}', \mathbf{p}) \end{pmatrix},$$

be L -operator in (3.35), (3.33) with $\mathbf{p}', \mathbf{p}$ in (3.25). Denote

$$\mathbf{L}_{m \sigma}^{m' \sigma''}(t; \mathbf{p}', \mathbf{p}) = \langle \langle \sigma | \mathbf{L}_m^{m'}(t; \mathbf{p}', \mathbf{p}) | \sigma'' \rangle \rangle, \quad \mathbf{L}_{m \sigma}^{\dagger m' \sigma''}(t; \mathbf{p}', \mathbf{p}) = \langle \langle \sigma | \mathbf{L}_m^{\dagger m'}(t; \mathbf{p}', \mathbf{p}) | \sigma'' \rangle \rangle,$$

where $\sigma, \sigma'' \in \mathbf{Z}_{\mathbf{n}}$, $m, m' = 0, 1$. Define

$$\begin{aligned} \mathbf{E}(\mathbf{p}')_{m; \sigma}^{\sigma''} &:= q^{-m \sigma} \mathcal{F}_{\mathbf{p}'}(\frac{-\sigma + \sigma''}{2}, m), & \mathbf{E}(\mathbf{p})_{\sigma}^{\sigma''; m'} &:= q^{m' \sigma''} \frac{\eta_{m'}}{\eta_{m'}} \mathcal{F}_{\mathbf{p}}(\frac{-\sigma + \sigma''}{2}, m'); \\ \mathbf{E}^\dagger(\mathbf{p}')_{m; \sigma}^{\sigma''} &:= \omega^{-m \sigma} F_{\mathbf{p}'}(\sigma - \sigma'', m), & \mathbf{E}^\dagger(\mathbf{p})_{\sigma}^{\sigma''; m'} &:= \omega^{m' \sigma''} \frac{\eta_{\sigma - \sigma''}}{\eta_{m'}} F_{\mathbf{p}}(\sigma - \sigma'', m'); \\ U_{\mathbf{p}, \mathbf{p}'}(d|_b^c) &:= \sum_{m=0,1} \mathbf{E}(\mathbf{p})_a^{d; m} \mathbf{E}(\mathbf{p}')_{m; b}^c, & U_{\mathbf{p}, \mathbf{p}'}^{\dagger}(d|_b^c) &:= \sum_{m=0,1} \mathbf{E}^\dagger(\mathbf{p})_a^{d; m} \mathbf{E}^\dagger(\mathbf{p}')_{m; b}^c \end{aligned} \quad (4.10)$$

for $a, b, c, d \in \mathbf{Z}_{\mathbf{n}}$ with $\frac{\eta_\sigma}{\eta_m}$ and $F_{\mathbf{p}}(\alpha, m)$ in (4.5). Then

(i) The (m, m') th entry of $\mathbf{L}, \mathbf{L}^\dagger$ -operator for $m, m' = 0, 1$ are expressed by

$$\mathbf{L}_{m \sigma}^{m' \sigma''}(\omega t; \mathbf{p}', \mathbf{p}) = \mathbf{E}(\mathbf{p}')_{m; \sigma}^{\sigma''} \mathbf{E}(\mathbf{p})_{\sigma}^{\sigma''; m'}, \quad \mathbf{L}_{m \sigma}^{\dagger m' \sigma''}(\omega t; \mathbf{p}', \mathbf{p}) = \mathbf{E}^\dagger(\mathbf{p}')_{m; \sigma}^{\sigma''} \mathbf{E}^\dagger(\mathbf{p})_{\sigma}^{\sigma''; m'},$$

where $\sigma, \sigma'' \in \mathbf{Z}_n$.

(ii) $U_{\mathbf{p}, \mathbf{p}'}$ and $U_{\mathbf{p}, \mathbf{p}'}^\dagger$ are expressed by

$$\begin{aligned} U_{\mathbf{p}, \mathbf{p}'}(d|c) &= \begin{cases} \langle \widehat{k^*} | L_n^{\dagger n'}(\omega t; \mathbf{p}^*, \mathbf{p}'^*) | \widehat{k^{*''}} \rangle & \text{if } (a-d, b-c) = (-2n, -2n'), n, n' = 0, 1; \\ 0 & \text{otherwise.} \end{cases} \\ U_{\mathbf{p}, \mathbf{p}'}^\dagger(d|c) &= \begin{cases} \langle \widehat{k^*} | L_n^{n'}(\omega t; \mathbf{p}^*, \mathbf{p}'^*) | \widehat{k^{*''}} \rangle & \text{if } (a-d, b-c) = (n, n'), n, n' = 0, 1; \\ 0 & \text{otherwise.} \end{cases} \end{aligned} \quad (4.11)$$

where $a-b=k^*, d-c=k^{*''} \in \mathbf{Z}_n$, and $\mathbf{p}^*, \mathbf{p}'^*$ are the dual of $\mathbf{p}, \mathbf{p}'$ in (4.3).

Proof. By the definition of $E(\mathbf{p}')$, $E(\mathbf{p})$ and $E^\dagger(\mathbf{p}')$, $E^\dagger(\mathbf{p})$ in (4.10), the spin-operator expressions of (3.29) (3.33) yield relations in (i). (ii) follows from the expression of $U_{\mathbf{p}, \mathbf{p}'}$, $U_{\mathbf{p}, \mathbf{p}'}^\dagger$ in (4.10):

$$\begin{aligned} U_{\mathbf{p}, \mathbf{p}'}(d|c) &= \sum_{m=0,1} q^{m(d-b)} (-\omega t)^{\frac{-a+d}{2}-m} \mathcal{F}_{\mathbf{p}}(\frac{-a+d}{2}, m) \mathcal{F}_{\mathbf{p}'}(\frac{-b+c}{2}, m), \\ U_{\mathbf{p}, \mathbf{p}'}^\dagger(d|c) &= \sum_{m=0,1} \omega^{m(d-b)} (-\omega t)^{a-d-m} F_{\mathbf{p}}(a-d, m) F_{\mathbf{p}'}(b-c, m) \end{aligned}$$

whose non-zero values are entries of the following respective L -operator:

$$\begin{aligned} &\begin{pmatrix} 1 - \omega t \frac{1}{bb'} \widehat{Z}' & (\frac{d'}{b'} - \omega \frac{a'd'}{bb'} \widehat{Z}') \widehat{X}'^2 \\ -\omega t (\frac{d}{b} - \frac{ad}{bb'} \widehat{Z}') \widehat{X}'^{-2} & -\omega t \frac{dd'}{bb'} + \omega \frac{aa'dd'}{bb'} \widehat{Z}' \end{pmatrix} = L^\dagger(\omega t; \mathbf{p}^*, \mathbf{p}'^*); \\ &\begin{pmatrix} 1 - \omega t \frac{1}{bb'} \widehat{Z}'^{-2} & (\frac{d'}{b'} - \omega \frac{a'd'}{bb'} \widehat{Z}'^{-2}) \widehat{X}'^{-1} \\ -\omega t (\frac{d}{b} - \frac{ad}{bb'} \widehat{Z}'^{-2}) \widehat{X}' & -\omega t \frac{dd'}{bb'} + \omega \frac{aa'dd'}{bb'} \widehat{Z}'^{-2} \end{pmatrix} = L(\omega t; \mathbf{p}^*, \mathbf{p}'^*). \end{aligned}$$

□

Using Lemma 4.2 (i) and (4.10), one finds the product form of $T^{(2)}(t; \mathbf{p}', \mathbf{p})$, $T^{\dagger(2)}(t; \mathbf{p}', \mathbf{p})$:

$$T^{(2)}(t; \mathbf{p}', \mathbf{p})_{\{\sigma_\ell\}}^{\{\sigma_\ell''\}} = \prod_{\ell=1}^L U_{\mathbf{p}, \mathbf{p}'}(\sigma_\ell'' | \sigma_{\ell+1}''), \quad T^{\dagger(2)}(t; \mathbf{p}', \mathbf{p})_{\{\sigma_\ell\}}^{\{\sigma_\ell''\}} = \prod_{\ell=1}^L U_{\mathbf{p}, \mathbf{p}'}^\dagger(\sigma_\ell'' | \sigma_{\ell+1}'').$$

By the boundary condition (3.42), one finds $E(\mathbf{p}')_m; \frac{\sigma_{L+1}''}{\sigma_{L+1}'} = q^{-mr'} E(\mathbf{p}')_m; \frac{\sigma_1''}{\sigma_1'}$, $E^\dagger(\mathbf{p}')_m; \frac{\sigma_{L+1}''}{\sigma_{L+1}'} = \omega^{-mr'} E^\dagger(\mathbf{p}')_m; \frac{\sigma_1''}{\sigma_1'}$, which provide the factor of the second term in (3.41) or (3.44). By (4.11), one finds

$$\begin{aligned} \langle \langle Q'; n_1, \dots, n_L | T^{(2)}(t; \mathbf{p}', \mathbf{p}) | Q'; n'_1, \dots, n'_L \rangle \rangle &= \langle \langle \widehat{n}_1, \dots, \widehat{n}_L | T^{\dagger(2)}(t; \mathbf{p}^*, \mathbf{p}'^*) | \widehat{n}'_1, \dots, \widehat{n}'_L \rangle \rangle; \\ \langle \langle Q'; n_1, \dots, n_L | T^{\dagger(2)}(t; \mathbf{p}', \mathbf{p}) | Q'; n'_1, \dots, n'_L \rangle \rangle &= \langle \langle \widehat{n}_1, \dots, \widehat{n}_L | T^{(2)}(t; \mathbf{p}^*, \mathbf{p}'^*) | \widehat{n}'_1, \dots, \widehat{n}'_L \rangle \rangle. \end{aligned}$$

Hence we obtain the duality between $\tau^{(2)}$ -models in (3.41), (3.44) as $\tau^{(2)}(t; \{\mathbf{p}'_{i_\ell}\}, \{\mathbf{p}_{i_\ell}\})$ in Proposition 4.1:

Proposition 4.2 *Let $\mathbf{p}^*, \mathbf{p}'^*$ be the dual of $\mathbf{p}, \mathbf{p}'$ in (4.3), and Ψ' be the dual correspondence between $V'_{r', Q'}$ and $V'_{r'^*, Q'^*}$ with $(r'^*, Q'^*) = (-Q, -r)$ in (4.9). Then $T^{(2)}(t; \mathbf{p}', \mathbf{p})$, $T^{\dagger(2)}(t; \mathbf{p}', \mathbf{p})$ on $V'_{r', Q'}$ is equivalent to $T^{\dagger(2)}(t; \mathbf{p}^*, \mathbf{p}'^*)$, $T^{(2)}(t; \mathbf{p}^*, \mathbf{p}'^*)$ on $V'_{r'^*, Q'^*}$ respectively by*

$$T^{(2)}(t; \mathbf{p}', \mathbf{p}) = \Psi'^{-1} T^{\dagger(2)}(t; \mathbf{p}^*, \mathbf{p}'^*) \Psi', \quad T^{\dagger(2)}(t; \mathbf{p}', \mathbf{p}) = \Psi'^{-1} T^{(2)}(t; \mathbf{p}^*, \mathbf{p}'^*) \Psi'. \quad (4.12)$$

□

4.3 Comparison of $\tau^{(2)}$ -duality and $t^{(2)}$ -duality

By Proposition 3.1 and 3.3, $T^{(2)}(t)$ and $T^{\dagger(2)}(t)$ can be decomposed as a sum of $\tau^{(2)}$ -models via the subspaces $\mathcal{C}^{\vec{i}}, \mathcal{C}^{\dagger\vec{i}}$ of $\bigotimes_{\ell=1}^L \mathbf{C}^{\mathbf{n}}$ for $\vec{i} \in \mathbf{Z}_2^L$ in (3.45) (3.48) respectively. Indeed, by (3.47), $T^{(2)}(t)|_{\mathcal{C}^{\vec{i}}} \simeq \tau^{(2)}(t; \{\mathbf{p}'_{i_\ell-1}\}, \{\mathbf{p}_{i_\ell}\})$ with $(\mathbf{p}'_{i_\ell}, \mathbf{p}_{i_\ell})$ in (3.46) and the boundary condition $r' \equiv 2r \pmod{\mathbf{n}}$. Similarly, $T^{\dagger(2)}(t)|_{\mathcal{C}^{\dagger\vec{i}}} \simeq \tau^{\dagger(2)}(t; \{\mathbf{p}'_{i_\ell}\}, \{\mathbf{p}_{i_\ell}\})$ with $(\mathbf{p}'_{i_\ell}, \mathbf{p}_{i_\ell})$ in (3.48) and the boundary condition $r' \equiv -r \pmod{N}$. One can lift the basis of $V_{r,Q}$ in (4.1) to $\mathcal{C}_{r,Q}^{\vec{i}}$ and $\mathcal{C}_{r,Q}^{\dagger\vec{i}}$, denoted by

$$\begin{aligned} \mathcal{C}_{r,Q}^{\vec{i}} &= \bigoplus_{k_\ell} \mathbf{C}|\widehat{k}_1, \dots, \widehat{k}_L\rangle\rangle_{\vec{i}} = \bigoplus_{n_\ell} \mathbf{C}|Q; n_1, \dots, n_L\rangle\rangle_{\vec{i}}, \\ \mathcal{C}_{r,Q}^{\dagger\vec{i}} &= \bigoplus_{k_\ell} \mathbf{C}|\widehat{k}_1, \dots, \widehat{k}_L\rangle\rangle_{\vec{i}}^\dagger = \bigoplus_{n_\ell} \mathbf{C}|Q; n_1, \dots, n_L\rangle\rangle_{\vec{i}}^\dagger, \end{aligned} \quad (4.13)$$

with $\sum_{\ell=1}^L k_\ell \equiv Q, \sum_{\ell=1}^L n_\ell \equiv r \pmod{N}$, where $|\widehat{k}_1, \dots, \widehat{k}_L\rangle\rangle_{\vec{i}}, |Q; n_1, \dots, n_L\rangle\rangle_{\vec{i}}, |\widehat{k}_1, \dots, \widehat{k}_L\rangle\rangle_{\vec{i}}^\dagger, |Q; n_1, \dots, n_L\rangle\rangle_{\vec{i}}^\dagger$ are basis elements corresponding to $|\widehat{k}_1, \dots, \widehat{k}_L\rangle, |Q; n_1, \dots, n_L\rangle$ of $V_{r,Q}$ via the projection $\wp_{\vec{i}}, \wp_{\vec{i}}^\dagger$ in (3.45) (3.48). By the expression of $\mathcal{C}^\pm, \mathcal{C}^{\dagger\pm}$ -basis in (3.3) and (3.11), one finds the relationship between the subspaces in (4.13) and (4.8) as follows:

$$\begin{aligned} \mathcal{C}_{r,Q}^{\vec{i}} &\subseteq V'_{r'Q'}, \quad \text{only if } Q = Q', 2r = r'; \\ \mathcal{C}_{r,Q}^{\dagger\vec{i}} &\subseteq V'_{r'Q'}, \quad \text{only if } i_0 = 0 : r' = -r, -2Q + \sum_{\ell} i_\ell \equiv Q', \end{aligned} \quad (4.14)$$

in particular, $\mathcal{C}_{r,Q}^{\dagger\vec{1}}$ is not contained in any $V'_{r'Q'}$. Hence in $\mathbf{n} = 2N$ case, the $T^{(2)}$ -duality in Proposition 4.2, with the $\tau^{(2)}$ -decomposition of $T^{(2)}(t), T^{\dagger(2)}(t)$ in Proposition 3.1, 3.3, is different from the $\tau^{(2)}$ -duality in Proposition 4.1. Indeed, the difference can also be seen in the inconsistency of parameter of $\tau^{(2)}$ -duality in Proposition 4.2:

$$\begin{aligned} T^{(2)}(t; \mathbf{p}', \mathbf{p})|_{\mathcal{C}^{\vec{1}}} &\simeq \tau^{(2)}(t; \{\mathbf{p}'_-\}, \{\mathbf{p}_-\}), & T^{\dagger(2)}(t; \mathbf{p}^*, \mathbf{p}'^*)|_{\mathcal{C}^{\dagger\vec{1}}} &\simeq \tau^{\dagger(2)}(t; \{\mathbf{p}^{*\dagger}_-\}, \{\mathbf{p}'^{*\dagger}_-\}); \\ T^{\dagger(2)}(t; \mathbf{p}', \mathbf{p})|_{\mathcal{C}^{\dagger\vec{1}}} &\simeq \tau^{\dagger(2)}(t; \{\mathbf{p}'^\dagger_-\}, \{\mathbf{p}^\dagger_-\}), & T^{(2)}(t; \mathbf{p}^*, \mathbf{p}'^*)|_{\mathcal{C}^{\vec{1}}} &\simeq \tau^{(2)}(t; \{\mathbf{p}^*_-\}, \{\mathbf{p}'^*_-\}). \end{aligned}$$

In $\mathbf{n} = N$ odd case, one can identify $\bigotimes_{\ell=1}^L \mathbf{C}^{\mathbf{n}}$ with $\mathcal{C}^{\vec{0}}$ or $\mathcal{C}^{\dagger\vec{0}}$ in Proposition 3.1 or 3.3.

Proposition 4.3 *When $\mathbf{n} = N$ odd, one has the identical quantum subspaces, $V'_{r'Q'} = \mathcal{C}_{r,Q}^{\vec{0}} = \mathcal{C}_{r^\dagger,Q^\dagger}^{\dagger\vec{0}}$ with $Q' \equiv Q \equiv -2Q^\dagger, r' \equiv 2r \equiv -r^\dagger \pmod{N}$, and the identification of basis elements in (4.8), (4.13):*

$$\begin{aligned} |\widehat{k}_1, \dots, \widehat{k}_L\rangle\rangle_{\vec{0}} &= |\widehat{k}_1, \dots, \widehat{k}_L\rangle\rangle, & |Q; n_1, \dots, n_L\rangle\rangle_{\vec{0}} &= |Q'; -2n_1, \dots, -2n_L\rangle\rangle, \\ |\widehat{k}_1^\dagger, \dots, \widehat{k}_L^\dagger\rangle\rangle_{\vec{0}}^\dagger &= |-\widehat{2k}_1^\dagger, \dots, -\widehat{2k}_L^\dagger\rangle\rangle^\dagger, & |Q^\dagger; n_1^\dagger, \dots, n_L^\dagger\rangle\rangle_{\vec{0}}^\dagger &= |Q'; n_1^\dagger, \dots, n_L^\dagger\rangle\rangle, \end{aligned} \quad (4.15)$$

with $\sum_{\ell=1}^L k_\ell \equiv Q, \sum_{\ell=1}^L k_\ell^\dagger \equiv Q^\dagger, \sum_{\ell=1}^L n_\ell \equiv r, \sum_{\ell=1}^L n_\ell^\dagger \equiv r^\dagger \pmod{N}$. With the identification of $\tau^{(2)}$ -models in Proposition 3.1 and 3.3,

$$\left(\tau^{(2)}(t; \{\mathbf{p}'\}, \{\mathbf{p}\}), \tau^{(2)}(t; \{\mathbf{p}^*\}, \{\mathbf{p}'^*\}) \right) \simeq \left(T^{(2)}(t; \mathbf{p}', \mathbf{p}), T^{\dagger(2)}(t; \mathbf{p}^*, \mathbf{p}'^*) \right) \simeq \left(T^{\dagger(2)}(t; \mathbf{p}', \mathbf{p}), T^{(2)}(t; \mathbf{p}^*, \mathbf{p}'^*) \right),$$

the duality relations in (4.12), (4.7) are the same.

Proof. By (3.3), (3.11) and (4.14), we obtain (4.15). On the dual space in (4.12), one has $V'_{r'^*Q'^*} = \mathcal{C}_{r^*, Q^*}^{\vec{0}} = \mathcal{C}_{r'^{\dagger}, Q'^{\dagger}}^{\dagger \vec{0}}$ with $Q'^* \equiv Q^* \equiv -2Q^{*\dagger}$, $r'^* \equiv 2r^* \equiv -r^{*\dagger}$, and the identification of basis elements:

$$\begin{aligned} |\widehat{k}_1^*, \dots, \widehat{k}_L^*\rangle_{\vec{0}} &= |\widehat{k}_1^*, \dots, \widehat{k}_L^*\rangle, & |Q^*; n_1^*, \dots, n_L^*\rangle_{\vec{0}} &= |Q'^*; -2n_1^*, \dots, -2n_L^*\rangle, \\ |\widehat{k}_1^{*\dagger}, \dots, \widehat{k}_L^{*\dagger}\rangle_{\vec{0}}^{\dagger} &= |-\widehat{2k}_1^{*\dagger}, \dots, -\widehat{2k}_L^{*\dagger}\rangle, & |Q^{*\dagger}; n_1^{*\dagger}, \dots, n_L^{*\dagger}\rangle_{\vec{0}}^{\dagger} &= |Q'^*; n_1^{*\dagger}, \dots, n_L^{*\dagger}\rangle, \end{aligned}$$

with $\sum_{\ell=1}^L k_{\ell}^* \equiv Q^*$, $\sum_{\ell=1}^L k_{\ell}^{*\dagger} \equiv Q^{*\dagger}$, $\sum_{\ell=1}^L n_{\ell}^* \equiv r^*$, $\sum_{\ell=1}^L n_{\ell}^{*\dagger} \equiv r^{*\dagger}$. Then the following conditions are equivalent,

$$(r'^*, Q'^*) = (-Q, -r) \Leftrightarrow (r^{*\dagger}, Q^{*\dagger}) = (Q, r) \Leftrightarrow (r^*, Q^*) = (Q^{\dagger}, r^{\dagger}).$$

Then the dual correspondence Ψ' in (4.9) becomes

$$|Q; n_1, \dots, n_L\rangle_{\vec{0}} \mapsto |n_1, \dots, n_L\rangle_{\vec{0}}^{\dagger}, \quad |Q^{\dagger}; n_1^{\dagger}, \dots, n_L^{\dagger}\rangle_{\vec{0}}^{\dagger} \mapsto |n_1^{\dagger}, \dots, n_L^{\dagger}\rangle_{\vec{0}},$$

by which, the dualities (4.12) and (4.7) are equivalent. $\square$

Other than the situation in Proposition 4.3, the $\tau^{(2)}$ -duality (4.7) in Proposition 4.1 can not be lifted to the duality between $T^{(2)}(t; \mathbf{p}', \mathbf{p})$ and $T^{\dagger(2)}(t; \mathbf{p}^*, \mathbf{p}'^*)$, neither between $T^{(2)}(t; \mathbf{p}', \mathbf{p})$ and $T^{(2)}(t; \mathbf{p}^*, \mathbf{p}'^*)$. In the latter case, the $\tau^{(2)}$ -model dual to (3.47) is $\tau^{(2)}(t; \{\mathbf{p}_{i_{\ell}}^*\}, \{\mathbf{p}_{i_{\ell+1}}'^*\})$, whose L -operator at ℓ th site in (3.39) is defined by the parameter

$$(\mathbf{p}_{i_{\ell}}^*, \mathbf{p}_{i_{\ell+1}}'^*) = (\mathbf{p}_+^*, \mathbf{p}_+'^*), (\mathbf{p}_-^*, \mathbf{p}_-'^*), (\mathbf{p}_+^*, \mathbf{p}_-'^*) \text{ or } (\mathbf{p}_-^*, \mathbf{p}_+'^*),$$

i.e. the l -twist (3.36) of $\mathbf{p}^*$ and $\mathbf{p}'^*$ with the identification $\mathbf{p}_{(-1)^l}^* = \mathbf{p}^*(l)$, $\mathbf{p}_{(-1)^l}'^* = \mathbf{p}'^*(-l)$ for $l = 0, 1$. Hence $\tau^{(2)}(t; \{\mathbf{p}_{i_{\ell}}^*\}, \{\mathbf{p}_{i_{\ell+1}}'^*\})$ is not a component of the decomposition of $T^{(2)}(t; \mathbf{p}^*, \mathbf{p}'^*)$ in Proposition 3.1. Since XXZ-model is related to $T^{(2)}$ -model by (3.43), the $\tau^{(2)}$ -duality in Proposition 4.1 can not be lifted to a duality among XXZ-models. Indeed, we shall shown in the next subsection, the dual model of XXZ-model $\mathcal{T}(s; \mathbf{p}', \mathbf{p})$ is given by $T^{(2)}(t; \mathbf{p}^*, \mathbf{p}'^*)$, so the essence of duality lies in the duality among $\tau^{(2)}$ -models.

4.4 Dual model of XXZ-model with cyclic representation

In this subsection, we derive the dual model of the XXZ-model $\mathcal{T}(s)$ in (3.41) with the boundary condition (3.42). As the L -operator of $T^{(2)}$ -model in Lemma 4.2, the L -operator of XXZ-model $\mathcal{T}(s)$ can be decomposed in the following product form:

Lemma 4.3 *Let*

$$\mathcal{L}(s; \mathbf{p}', \mathbf{p}) = \begin{pmatrix} \mathcal{L}_+^+(s; \mathbf{p}', \mathbf{p}) & \mathcal{L}_-^+(s; \mathbf{p}', \mathbf{p}) \\ \mathcal{L}_-^-(s; \mathbf{p}', \mathbf{p}) & \mathcal{L}_+^-(s; \mathbf{p}', \mathbf{p}) \end{pmatrix}, \quad L^{\dagger}(t; \mathbf{p}^*, \mathbf{p}'^*) = \begin{pmatrix} L_+^{\dagger+}(t; \mathbf{p}^*, \mathbf{p}'^*) & L_+^{\dagger-}(t; \mathbf{p}^*, \mathbf{p}'^*) \\ L_-^{\dagger+}(t; \mathbf{p}^*, \mathbf{p}'^*) & L_-^{\dagger-}(t; \mathbf{p}^*, \mathbf{p}'^*) \end{pmatrix}$$

be the L -operator in (3.29), (3.33) respectively, where $\mathbf{p}' = (\mathbf{a}', \mathbf{b}', \mathbf{d}')$, $\mathbf{p} = (\mathbf{a}, \mathbf{b}, \mathbf{d})$ and $\mathbf{p}^*, \mathbf{p}'^*$ are the dual of $\mathbf{p}, \mathbf{p}'$ in (4.3), (here we use $\gamma = \pm (= \pm 1)$ in (3.38) as the auxiliary index of the L -operator). Denote

$$\mathcal{L}_{\gamma}^{\gamma' \sigma''}(s) = \langle \langle \sigma | \mathcal{L}_{\gamma}^{\gamma'}(s; \mathbf{p}', \mathbf{p}) | \sigma'' \rangle \rangle, \quad L_{\delta}^{\dagger \delta' k''}(t) = \langle \langle \widehat{k}^* | \mathcal{L}_{\delta}^{\dagger \delta'}(t; \mathbf{p}^*, \mathbf{p}'^*) | \widehat{k}'' \rangle \rangle$$

where $\sigma, \sigma'', k^*, k^{*''} \in \mathbf{Z}_n$, $\gamma, \gamma', \delta, \delta' = \pm$. Define

$$\begin{aligned}\mathcal{E}(\mathbf{p}')_{\gamma; \sigma}^{\sigma''} &:= (\mathbf{b}'\mathbf{b})^{\frac{m}{2}} q^{-m\sigma} (-s)^{-(1+m)} \mathcal{F}_{\mathbf{p}'}(\sigma - \sigma'', \gamma), \quad (m = \frac{1-\gamma}{2}), \\ \mathcal{E}(\mathbf{p})_{\sigma}^{\sigma''; \gamma'} &:= (\mathbf{b}'\mathbf{b})^{\frac{-m'}{2}} q^{m'(\sigma''+1)} (-s)^{m'} \frac{\eta_{\sigma-\sigma''}}{\eta_{\gamma'}} \mathcal{F}_{\mathbf{p}}(\sigma - \sigma'', \gamma'), \quad (m' = \frac{1-\gamma'}{2}); \\ \mathcal{U}_{\mathbf{p}, \mathbf{p}'}(d|c) &:= \sum_{\gamma=\pm} \mathcal{E}(\mathbf{p})_a^d \gamma \mathcal{E}(\mathbf{p}')_{\gamma; b}^c,\end{aligned}\tag{4.16}$$

for $a, b, c, d \in \mathbf{Z}_n$, where $\frac{\eta_-}{\eta_+} = -s^2$, $\mathcal{F}_{\mathbf{p}}(+, +) = (\frac{q\mathbf{b}^3}{a d^2})^{\frac{1}{4}}$, $\mathcal{F}_{\mathbf{p}}(+, -) = (\frac{-s^2}{b})(\frac{q\mathbf{b}^3}{a d^2})^{\frac{1}{4}}$, $\mathcal{F}_{\mathbf{p}}(-, +) = (\frac{d}{b})(\frac{q\mathbf{b}^3}{a d^2})^{\frac{1}{4}}$, $\mathcal{F}_{\mathbf{p}}(-, -) = (\frac{-\omega a d}{b})(\frac{q\mathbf{b}^3}{a d^2})^{\frac{1}{4}}$, and $\mathcal{F}_{\mathbf{p}}(\alpha, \gamma) = 0$ if $\alpha \neq \pm 1$. Then

(i) $\mathcal{L}_{\gamma}^{\gamma' \sigma''}(s) = \mathcal{E}(\mathbf{p}')_{\gamma; \sigma}^{\sigma''} \mathcal{E}(\mathbf{p})_{\sigma}^{\sigma''; \gamma'}$ for $\sigma, \sigma'' \in \mathbf{Z}_n$, $\gamma, \gamma' = \pm$.

(ii) $\mathcal{U}_{\mathbf{p}, \mathbf{p}'}(d|c) = 0$ if $a - d$ or $b - c \neq \pm 1$, and

$$\mathcal{U}_{\mathbf{p}, \mathbf{p}'}(d|c) = -s^{-1} (\frac{q^2 \mathbf{b}^* \mathbf{b}'^*}{a^* a'^* c^{*2}})^{\frac{1}{4}} L_{\delta}^{\dagger \delta' k^* k^{*''}}(t) \quad \text{when } (a - d, b - c) = (\delta, \delta'), \delta, \delta' = \pm 1,$$

where $k^* = a - b, k^{*''} = d - c \in \mathbf{Z}_n$.

Proof. (i) follows from the spin-operator expression of $\mathcal{L}(s; \mathbf{p}', \mathbf{p})$ in (3.29) and the definition of $\mathcal{E}(\mathbf{p}')_{\gamma; \sigma}^{\sigma''}$ and $\mathcal{E}(\mathbf{p})_{\sigma}^{\sigma''; \gamma'}$. One also finds

$$\mathcal{U}_{\mathbf{p}, \mathbf{p}'}(d|c) = -s^{-1} \sum_{\gamma=\pm} q^{\frac{(1-\gamma)(d-b+1)}{2}} (-s^2)^{\frac{\gamma-a+d}{2}} \mathcal{F}_{\mathbf{p}}(a - d, \gamma) \mathcal{F}_{\mathbf{p}'}(b - c, \gamma),$$

of which the non-zero values form the following L -operator

$$-s^{-1} (\frac{q^2 \mathbf{b}^* \mathbf{b}'^*}{a a' c^2})^{\frac{1}{4}} \begin{pmatrix} 1 - t \frac{1}{b b'} \widehat{Z}' & (\frac{d'}{b'} - \omega \frac{a' d'}{b b'}) \widehat{X}'^2 \\ -t (\frac{d}{b} - \frac{a d}{b b'}) \widehat{Z}' \widehat{X}'^{-2} & -t \frac{d d'}{b b'} + \omega \frac{a a' d d'}{b b'} \widehat{Z}' \end{pmatrix} = -s^{-1} (\frac{q^2 \mathbf{b}^* \mathbf{b}'^*}{a^* a'^* c^{*2}})^{\frac{1}{4}} L^{\dagger}(t; \mathbf{p}^*, \mathbf{p}'^*).$$

Then (ii) follows. $\square$

By Lemma 4.3 (i), $\mathcal{T}(s; \mathbf{p}', \mathbf{p})$ can be expressed in the product form:

$$\mathcal{T}(s; \mathbf{p}', \mathbf{p})_{\{\sigma_{\ell}\}}^{\{\sigma_{\ell}''\}} = \prod_{\ell=1}^L \mathcal{U}_{\mathbf{p}, \mathbf{p}'}(\sigma_{\ell}'' | \sigma_{\ell+1}''),$$

with the relation $\mathcal{E}(\mathbf{p}')_{\gamma; \sigma_{L+1}''} = q^{-\frac{(1-\gamma)}{2} r'} \mathcal{E}(\mathbf{p}')_{\gamma; \sigma_1''}$ by (3.42), which contributes the $q^{-r'}$ -factor in (3.41). Then Lemma 4.3 (ii) yields the relation

$$\begin{aligned}& \langle \langle Q'; n_1, \dots, n_L | \mathcal{T}(s; \mathbf{p}', \mathbf{p}) | Q'; n'_1, \dots, n'_L \rangle \rangle \\ &= (-s)^{-L} (\frac{q^2 \mathbf{b}^* \mathbf{b}'^*}{a^* a'^* c^{*2}})^{\frac{L}{4}} \langle \langle \widehat{n}_1, \dots, \widehat{n}_L | T^{\dagger(2)}(\omega^{-1} t; \mathbf{p}^*, \mathbf{p}'^*) | \widehat{n}'_1, \dots, \widehat{n}'_L \rangle \rangle.\end{aligned}$$

Hence up to a scale function, $T^{\dagger(2)}(\omega^{-1} t; \mathbf{p}^*, \mathbf{p}'^*)$ is the dual model of $\mathcal{T}(s; \mathbf{p}', \mathbf{p})$:

Proposition 4.4 *Let $\mathbf{p}^*, \mathbf{p}'^*$ be the dual of $\mathbf{p}, \mathbf{p}'$ in (4.3), and Ψ' be the dual correspondence between $V_{r', Q'}$ and $V_{r'^*, Q'^*}$ with $(r'^*, Q'^*) = (-Q, -r)$ in (4.9). Then $\mathcal{T}(s; \mathbf{p}', \mathbf{p})$ on $V_{r', Q'}$ is equivalent to $(-s)^{-L} (\frac{q^2 \mathbf{b}^* \mathbf{b}'^*}{a^* a'^* c^{*2}})^{\frac{L}{4}} T^{\dagger(2)}(\omega^{-1} t; \mathbf{p}^*, \mathbf{p}'^*)$ on $V_{r'^*, Q'^*}$ by*

$$\mathcal{T}(s; \mathbf{p}', \mathbf{p}) = (-s)^{-L} (\frac{q^2 \mathbf{b}^* \mathbf{b}'^*}{a^* a'^* c^{*2}})^{\frac{L}{4}} \Psi'^{-1} T^{\dagger(2)}(\omega^{-1} t; \mathbf{p}^*, \mathbf{p}'^*) \Psi'. \tag{4.17}$$

□

By Proposition 4.2, $\mathsf{T}^{\dagger(2)}(t; \mathbf{p}', \mathbf{p})$ is also the dual model of $\mathsf{T}^{(2)}(t; \mathbf{p}', \mathbf{p})$, which is related to $\mathcal{T}(s; \mathbf{p}', \mathbf{p})$ by (3.43). Note that the duality in (4.6) or (4.12) is the identification of dual $\tau^{(2)}$ -transfer matrices, not as the ABCD-algebra representation in (2.8). In particular, the operator $\widehat{Z}'$ in (3.43) is not preserved under the duality correspondence Ψ' in (4.12). This explains the reason why $\mathsf{T}^{\dagger(2)}(t; \mathbf{p}', \mathbf{p})$ serves the dual model for both $\mathsf{T}^{(2)}(t; \mathbf{p}', \mathbf{p})$ and $\mathcal{T}(s; \mathbf{p}', \mathbf{p})$.

5 Inhomogeneous Chiral Potts Model and XXZ model with cyclic $U_q(sl_2)$ -representation

Let ω, q be roots of unity in (1.1). For convenience, we may assume $\omega = e^{\frac{2\pi i}{N}}$, and $q = -\omega^{\frac{-1}{2}}$ when $\mathbf{n} = N$ odd, and $q = \pm\omega^{\frac{-1}{2}}$ when $\mathbf{n} = 2N$, where $\omega^{\frac{1}{2}} := e^{\frac{\pi i}{N}}$. By the gauge equivalence and the scaling of spectral parameter, the five parameters in the L -operator of $\tau^{(2)}$ -model in (3.23) (3.25) can be reduced to the 3-parameter family represented by two rapidities in the same k' -curve of CPM model [24]:

$$(\mathbf{a}', \mathbf{b}', \mathbf{a}, \mathbf{b}, \mathbf{c}) = (x_{p'}, y_{p'}, x_p, y_p, \mu_{p'}\mu_p), \quad \mathbf{p}' = p' = (x_{p'}, y_{p'}, \mu_{p'}), \quad \mathbf{p} = p = (x_p, y_p, \mu_p), \quad (5.1)$$

with $p, p' \in \mathfrak{W}_{k'}$, defined by

$$\mathfrak{W}_{k'} (= \mathfrak{W}_{k', k}) : kx^N = 1 - k'\mu^{-N}, ky^N = 1 - k'\mu^N, (k'^2 \neq 0, 1, k^2 + k'^2 = 1) \quad (5.2)$$

(see, e.g. [1, 9]). Here, $\mathfrak{W}_{k'}$ is represented by $\mathfrak{W}_{k', k}$ or $\mathfrak{W}_{k', -k}$, which are isomorphic via the transformation $(x, y, \mu) \mapsto (\pm\omega^{\frac{1}{2}}x, \pm\omega^{\frac{1}{2}}y, \mu)$.

5.1 Duality in chiral Potts model

In chiral Potts model, the $\tau^{(2)}$ -model (3.39) are defined by the L -operators with the parameter $\mathbf{p}'_{i_\ell} = p'_{i_\ell}, \mathbf{p}_{i_\ell} = p_{i_\ell} \in \mathfrak{W}_{k'}$ in (5.2):

$$\tau^{(2)}(t) = \tau^{(2)}(t; \{p'_{i_\ell}\}, \{p_{i_\ell}\}), \quad t := x_q y_q \quad \text{for } q \in \mathfrak{W}_{k'}. \quad (5.3)$$

It is known that the Q -operator of $\tau^{(2)}$ -model (5.3) in the theory of Baxter's TQ -relation [5] is the $\overset{L}{\otimes} \mathbf{C}^N$ -transfer matrix in CPM [8, 9, 23, 24]³:

$$\begin{aligned} T(q)_{\{\sigma\}, \{\sigma'\}} &= T(q; \{p'_\ell\}, \{p_\ell\})_{\{\sigma\}, \{\sigma'\}} = \prod_{\ell=1}^L W_{p_\ell q}(\sigma_\ell - \sigma'_\ell) \overline{W}_{p'_{\ell+1} q}(\sigma_{\ell+1} - \sigma'_\ell), \\ \widehat{T}(q)_{\{\sigma'\}, \{\sigma''\}} &= \widehat{T}(q; \{p'_\ell\}, \{p_\ell\})_{\{\sigma'\}, \{\sigma''\}} = \prod_{\ell=1}^L \overline{W}_{p_\ell q}(\sigma'_\ell - \sigma''_\ell) W_{p'_{\ell+1} q}(\sigma'_\ell - \sigma''_{\ell+1}), \end{aligned} \quad (5.4)$$

where $q \in \mathfrak{W}_{k'}$ in (5.2), $\sigma, \sigma', \sigma'' \in \mathbf{Z}_N$ are the spin-basis in (1.2), and $W_{pq}, \overline{W}_{pq}$ are the Boltzmann weights in CPM [10]:

$$\frac{W_{pq}(\sigma)}{W_{pq}(0)} = \left(\frac{\mu_p}{\mu_q}\right)^\sigma \prod_{j=1}^\sigma \frac{y_q - \omega^j x_p}{y_p - \omega^j x_q}, \quad \frac{\overline{W}_{pq}(\sigma)}{\overline{W}_{pq}(0)} = (\mu_p \mu_q)^\sigma \prod_{j=1}^\sigma \frac{\omega x_p - \omega^j x_q}{y_q - \omega^j y_p} \quad (5.5)$$

³We shift the index of p'_ℓ in [27] (2.16) by one, where p'_ℓ is equal to $p'_{\ell+1}$ of (5.4) here.

with $W_{p,q}(0) = \overline{W}_{p,q}(0) = 1$. Here both (5.3) and (5.4) are with the boundary condition (3.40) and the periodic-vertical rapidities $(p'_{L+1}, p_{L+1}) = (p'_1, p_1)$. The star-triangle relation of Boltzmann weights [2, 3, 10, 18, 19] yields

$$T(q)\hat{T}(r) = \left(\frac{f_{p'q}f_{pr}}{f_{pq}f_{p'r}}\right)^L T(r)\hat{T}(q), \quad \hat{T}(q)T(r) = \left(\frac{f_{pq}f_{p'r}}{f_{p'q}f_{pr}}\right)^L \hat{T}(r)T(q),$$

with the commutative relations,

$$[\hat{T}(q)T(r), \hat{T}(q')T(r')] = [T(q)\hat{T}(r), T(q')\hat{T}(r')] = 0$$

for $q, r, q', r' \in \mathfrak{W}_{k'}$, where $f_{pq} = (\frac{\det_N(\overline{W}_{pq}(i-j))}{\prod_{n=0}^{N-1} W_{pq}(n)})^{1/N}$. Then $T(q), \hat{T}(q)$ can be diagonalized via two invertible q -independent matrices P_B, P_W , i.e. $P_W^{-1}T(q)P_B = T_{\text{diag}}(q)$, $P_B^{-1}\hat{T}(q)P_W = \hat{T}_{\text{diag}}(q)$ diagonal so that $\hat{T}_{\text{diag}}(q) = T_{\text{diag}}(q)(\frac{f_{pq}}{f_{p'q}})^L D$ for some q -independent diagonal matrix D ([9] (2.32)-(2.34), (4.46), [8] (2.10)-(2.13)). The Baxter's TQ -relation is in the form of $\tau^{(2)}T$ -relation between (5.3) and (5.4) ([8] (3.15), [9] (4.31), [25] (2.31)-(2.32)):

$$\begin{aligned} \tau^{(2)}(t_q)T(Uq) &= \varphi_q T(q) + \omega^r \overline{\varphi}_{Uq} XT(U^2q), \\ \tau^{(2)}(t_q)T(U'q) &= \omega^r \varphi'_q XT(q) + \overline{\varphi}'_{U'q} T(U'^2q), \end{aligned} \quad (5.6)$$

where U, U' are automorphisms of (5.2) defined by $U(x, y, \mu) = (\omega x, y, \mu)$, $U'(x, y, \mu) = (x, \omega y, \mu)$, and $\varphi_q (= \varphi_{\{p'_\ell\}, \{p_\ell\}; q})$, $\overline{\varphi}_q (= \overline{\varphi}_{\{p'_\ell\}, \{p_\ell\}; q})$, $\varphi'_q (= \varphi'_{\{p'_\ell\}, \{p_\ell\}; q})$, $\overline{\varphi}'_q (= \overline{\varphi}'_{\{p'_\ell\}, \{p_\ell\}; q})$ are functions defined by

$$\begin{aligned} \varphi_q &= \prod_\ell \frac{(t_{p'_\ell} - t_q)(y_{p_\ell} - \omega x_q)}{y_{p_\ell} y_{p'_\ell} (x_{p'_\ell} - x_q)}, & \overline{\varphi}_q &= \prod_\ell \frac{\omega \mu_{p'_\ell} \mu_{p_\ell} (t_{p_\ell} - t_q)(x_{p'_\ell} - x_q)}{y_{p_\ell} y_{p'_\ell} (y_{p_\ell} - \omega x_q)}, \\ \varphi'_q &= \prod_\ell \frac{\omega \mu_{p_\ell} \mu_{p'_\ell} (t_{p'_\ell} - t_q)(x_{p_\ell} - y_q)}{y_{p_\ell} y_{p'_\ell} (y_{p'_\ell} - y_q)}, & \overline{\varphi}'_q &= \prod_\ell \frac{(t_{p_\ell} - t_q)(y_{p'_\ell} - y_q)}{y_{p_\ell} y_{p'_\ell} (x_{p_\ell} - y_q)}. \end{aligned}$$

We now discuss the duality of $\tau^{(2)}$ -model (5.3) in CPM. In order to preserve the vertical rapidities in (5.3), we modify the parameter-duality (4.3) by a factor $\alpha = i^{\frac{1}{N}}$ in the Remark of Proposition 4.1, so that the parameter-duality defines the dual rapidities between k' - and k'^{-1} -curves $\mathfrak{W}_{k'}$ and $\mathfrak{W}_{1/k'}$ ([27] (3.9)):

$$\mathfrak{W}_{k'} \xrightarrow{\sim} \mathfrak{W}_{1/k'}, \quad p = (x_p, y_p, \mu_p) \mapsto p^* = (x_{p^*}, y_{p^*}, \mu_{p^*}) := (i^{\frac{1}{N}} x_p \mu_p, i^{\frac{1}{N}} y_p \mu_p^{-1}, \mu_p^{-1}). \quad (5.7)$$

Then the $\tau^{(2)}$ -duality (4.7) takes the following form:

$$\tau^{(2)}(t; \{p'_{i_\ell}\}, \{p_{i_\ell}\}) = \Psi^{-1} \tau^{(2)}(t^*; \{p^*_{i_\ell}\}, \{p'^*_{i_\ell+1}\}) \Psi, \quad t^* = (-1)^{\frac{1}{N}} t. \quad (5.8)$$

The duality (5.8) can be extended to the duality of CPM. Indeed, $T, \hat{T}$ in (5.4) commute with the charge operator $\hat{Z}(= X)$, hence preserve the Q -subspace $V_{r,Q}$ in (4.1). The Boltzmann weights (5.5) for dual rapidities in (5.7) are related by $\frac{\overline{W}_{pq}^{(f)}(k)}{\overline{W}_{pq}^{(f)}(0)} = W_{p^*q^*}(k)$, $\frac{W_{pq}^{(f)}(k)}{W_{pq}^{(f)}(0)} = \overline{W}_{p^*q^*}(N-k)$, where $\overline{W}_{pq}^{(f)}(k) = \frac{1}{\sqrt{N}} \sum_{\sigma=0}^{N-1} \omega^{k\sigma} \overline{W}_{pq}(\sigma)$, $W_{pq}^{(f)}(k) = \frac{1}{\sqrt{N}} \sum_{\sigma=0}^{N-1} \omega^{k\sigma} W_{pq}(\sigma)$ are the Fourier transform of Boltzmann weights in (5.5), ([27] (3.17)). By [27] Theorem 3.1, when $(r^*, Q^*) = (Q, r)$, the

chiral Potts transfer matrices (5.4), $T(q; \{p'_\ell\}, \{p_\ell\})$, $\widehat{T}(q; \{p'_\ell\}, \{p_\ell\})$ on $V_{r,Q}$ and $T(q^*; \{p_\ell^*\}, \{p'_{\ell+1}\})$, $\widehat{T}(q^*; \{p_\ell^*\}, \{p'_{\ell+1}\})$ on V_{r^*,Q^*} , are equivalent via the dual correspondence Ψ in (4.2):

$$\begin{aligned} T(q^*; \{p_\ell^*\}, \{p'_{\ell+1}\}) &= \left(\prod_\ell \frac{W_{p'_\ell, q^*}^{(f)}(0)}{W_{p_\ell, q}^{(f)}(0)} \right) \Psi T(q; \{p'_\ell\}, \{p_\ell\}) \Psi^{-1}, \\ \widehat{T}(q^*; \{p_\ell^*\}, \{p'_{\ell+1}\}) &= \left(\prod_\ell \frac{W_{p'_\ell, q^*}^{(f)}(0)}{W_{p_\ell, q}^{(f)}(0)} \right) \Psi \widehat{T}(q; \{p'_\ell\}, \{p_\ell\}), \end{aligned} \quad (5.9)$$

where $p_\ell^*, p'_{\ell+1}, q^* \in \mathfrak{W}_{1/k'}^*$ are the dual of $p_\ell, p'_\ell, q \in \mathfrak{W}_{k'}$ in (5.7).

5.2 Chiral Potts transfer matrix as the Q -operator of XXZ-model with cyclic $U_q(sl_2)$ -representation

In this subsection, we study the $T^{(2)}$ - and XXZ-model, (3.41) and (3.44) with parameter in (5.1): $p' = p', p = p \in \mathfrak{W}_{k'}$. Then (p'_-, p_-) in (3.27) and $(p_-^\dagger, p_-^\dagger)$ in (3.48) become

$$p'_- = (q^{-1}x_{p'}, qy_{p'}, \mu_{p'}), \quad p_- = (qx_p, q^{-1}y_p, \mu_p), \quad p_-^\dagger = p', \quad p_-^\dagger = (x_p, y_p, q\mu_p),$$

which are elements in $\mathfrak{W}_{k'}$ in $\mathbf{n} = N$ odd case, denoted by $p'_-, p_-, p_-^\dagger$. In $\mathbf{n} = 2N$ case, p'_-, p_- are elements in $\mathfrak{W}_{k', -k}$, identified with $p'_- = (x_{p'}, \omega^{-1}y_{p'}, \mu_{p'}), p_- = (\omega^{-1}x_p, y_p, \mu_p) \in \mathfrak{W}_{k'} (= \mathfrak{W}_{k', k})$; and $p_-^\dagger = p_-^\dagger \in \mathfrak{W}_{-k'}$. Hence parameters in (3.27) and (3.48) are identified with

$$\begin{aligned} \mathbf{n} = N \text{ odd : } & p'_- = (q^{-1}x_{p'}, qy_{p'}, \mu_{p'}), p_- = (qx_p, q^{-1}y_p, \mu_p), p_-^\dagger = p', p_-^\dagger = (x_p, y_p, q\mu_p), \\ & (p'_-, p_-, p_-^\dagger, p_-^\dagger) \in \mathfrak{W}_{k'}; \\ \mathbf{n} = 2N : & p'_- = (x_{p'}, \omega^{-1}y_{p'}, \mu_{p'}), p_- = (\omega^{-1}x_p, y_p, \mu_p), p_-^\dagger = p', p_-^\dagger = (x_p, y_p, q\mu_p), \\ & (p'_-, p_-, p_-^\dagger, p_-^\dagger) \in \mathfrak{W}_{k'}, p_-^\dagger \in \mathfrak{W}_{-k'}. \end{aligned} \quad (5.10)$$

Over the subspace $\mathcal{C}^{\vec{i}}$ of $\bigotimes^L \mathbf{C}^{\mathbf{n}}$ in (3.45) for $\vec{i} \in \mathbf{Z}_2^L$, the equivalence of $T^{(2)}(t) (= T^{(2)}(t; p', p))$ and $\tau^{(2)}$ -model in (3.47) becomes

$$T^{(2)}(t)|_{\mathcal{C}^{\vec{i}}} \simeq \tau^{(2)}(t; \{p'_{i_\ell}\}, \{p_{i_\ell}\}), \quad p'_{i_\ell} = p'_{(-1)^{i_\ell}}, p_{i_\ell} = p_{(-1)^{i_\ell}} \in \mathfrak{W}_{k'}. \quad (5.11)$$

Hence we may regard $T(q; \{p'_\ell\}, \{p_\ell\})$, $\widehat{T}(q; \{p'_\ell\}, \{p_\ell\})$ in (5.4) as the Q -operator of $T^{(2)}(t)|_{\mathcal{C}^{\vec{i}}}$, with $\tau^{(2)}T$ -relation induced from (5.6). In $\mathbf{n} = N$ odd case, by Proposition 3.1 (i), $\bigotimes^L \mathbf{C}^{\mathbf{n}} = \mathcal{C}^{\vec{i}}$; then by Proposition 3.2 (i), the XXZ-model $\mathcal{T}(s, p', p)$ in (3.41) can be identified with $T^{(2)}(t)$ up to some scale function, by which one may consider $T(q; \{p'_\ell\}, \{p_\ell\})$, $\widehat{T}(q; \{p'_\ell\}, \{p_\ell\})$ as the Q -operator of $\mathcal{T}(s, p', p)$. On the other hand, since $p_-^\dagger, p_-^\dagger \in \mathfrak{W}_{k'}$ by (5.10), the Q -operators of the equivalent models in Proposition 3.3 (i), $T^{\dagger(2)}(t)|_{\mathcal{C}^{\dagger\vec{i}}} \simeq \tau^{\dagger(2)}(t; \{p_{i_\ell}^\dagger\}, \{p_{i_\ell}^\dagger\})$ with $p_{i_\ell}^\dagger = p_{(-1)^{i_\ell}}^\dagger, p_{i_\ell}^\dagger = p_{(-1)^{i_\ell}}^\dagger \in \mathfrak{W}_{k'}$, are $T(q; \{p_{i_\ell}^\dagger\}, \{p_{i_\ell}^\dagger\})$, $\widehat{T}(q; \{p_{i_\ell}^\dagger\}, \{p_{i_\ell}^\dagger\})$. In particular, when $\vec{i} = \vec{0}$ where $p'_{i_\ell} = p_{i_\ell}^\dagger = p', p_{i_\ell} = p_{i_\ell}^\dagger = p$ for all ℓ , the duality in Proposition 4.3 extends to the duality of Q -operators with $p'_\ell = p', p_\ell = p, p_\ell^* = p'^*, p_\ell^* = p^*$ in (5.9).

In $\mathbf{n} = 2N$ case, since $p_-^\dagger \in \mathfrak{W}_{k'}, p_-^\dagger \in \mathfrak{W}_{-k'}$ in (5.10), the vertical rapidities $p_{i_\ell}^\dagger, p_{i_\ell}^\dagger$ of the $\tau^{(2)}$ -model equivalent to $T^{\dagger(2)}(t)$ on the component $\mathcal{C}^{\dagger\vec{i}}$ in Proposition 3.3 (ii) are not all in the

same k' -curve when $\vec{i} \neq \vec{0}$, hence the chiral Potts transfer matrix fails to be the Q -operator of $T^{(2)}$ -model. For $T^{(2)}$ -model in Proposition 3.1 (ii), since the parameter ν in (3.24) corresponding to (p'_-, p_-) in (3.27) and (p'_-, p_-) in (5.10) differ by ω in $\mathbf{n} = 2N$ case, $\tau^{(2)}(t; \{p'_{i_\ell}\}, \{p_{i_\ell}\})$ in (5.11) and $\tau^{(2)}(t; \{p'_{i_\ell}\}, \{p_{i_\ell}\})$ in (3.47) are related by the relation (2.13) with $\xi_\ell = \omega^{i_\ell}$. Therefore, the $\widehat{Z}'$ -identification between $T^{(2)}(t)_{|C^{\vec{i}}}$ and $T^{(2)}(t)_{|C^{\vec{i}+\vec{1}}}$ in Proposition 3.2 (ii) corresponds the identification of $\tau^{(2)}(t; \{p'_{i_\ell}\}, \{p_{i_\ell}\})$ and $\tau^{(2)}(t; \{p'_{i_\ell+1}\}, \{p_{i_\ell+1}\})$ in (5.11) via the change of local spectral parameters at site ℓ , $t \mapsto \omega^{(-1)^{i_\ell+1}} t$. Then $T(q; \{p'_\ell\}, \{p_\ell\}), \widehat{T}(q; \{p'_\ell\}, \{p_\ell\})$ are identified with $T(q; \{p'_\ell+1\}, \{p_\ell+1\}), \widehat{T}(q; \{p'_\ell+1\}, \{p_\ell+1\})$ with the compatible $\tau^{(2)}T$ -relation (5.6) via the change of variable $q \in \mathfrak{W}_{k'}$ at site ℓ : $(x_q, y_q, \mu_q) \mapsto (x_q, \omega^{\mp 1} y_q, \mu_q)$ according to $(p'_{i_\ell} p_{i_\ell}) = (p'_\pm, p_\pm)$. Therefore, the two pairs of chiral Potts transfer matrices, $(T(q; \{p'_\ell\}, \{p_\ell\}), T(q; \{p'_\ell+1\}, \{p_\ell+1\}))$ and $(\widehat{T}(q; \{p'_\ell\}, \{p_\ell\}), \widehat{T}(q; \{p'_\ell+1\}, \{p_\ell+1\}))$, form the Q -operator of $\mathcal{T}(s; p', p)$ over the component $\mathcal{C}^{[i]}$ of $\bigotimes^L \mathbf{C}^{\mathbf{n}}$ in Proposition 3.2 (ii).

6 Concluding Remarks

In this paper, we first in section 2 characterize the quantum group $U_w(sl_2)$ in the L -operator of $\tau^{(2)}$ -model for a generic w as a subalgebra of the quantum group $U_q(sl_2)$ in XXZ model with $q^{-2} = w$. Then we study the XXZ model with cyclic representation of $U_q(sl_2)$, and its related $\tau^{(2)}$ -models in the root of unity case (1.1). Through the representation theory of $U_\omega(sl_2)$, we obtain the structure of XXZ- and $\tau^{(2)}$ -models, and their relationship with non-superintegrable N -state CPM in sections 3 and 5. We also study the duality of XXZ- and $\tau^{(2)}$ -models with cyclic representation in section 4, and find the fundamental role of $\tau^{(2)}$ -models in the duality theory. One expects a similar structure will appear again in XXZ models defined by other representations of $U_q(sl_2)$, not only for the cyclic representation in this work. The representation theory should provide a more direct access to the eigenvector problem of models related to $\tau^{(2)}$ -model, as indicated in [28] about the eigenvectors of superintegrable chiral Potts model. A program along this line is now under consideration, and progress is expected. We leave the discussion to future work.

References

- [1] G. Albertini, B. M. McCoy, and J. H. H. Perk, Eigenvalue spectrum of the superintegrable chiral Potts model, In *Integrable system in quantum field theory and statistical mechanics*, Adv. Stud. Pure Math., 19, Kinokuniya Academic, Academic Press, Boston, MA (1989) 1–55.
- [2] H. Au-Yang, B. M. McCoy, J. H. H. Perk and S. Tang, Solvable models in statistical mechanics and Riemann surfaces of genus greater than one, *Algebraic Analysis*, Vol. 1, eds. M. Kashiwara and T. Kawai, Academic Press, San Diego (1988), 29–40.
- [3] H. Au-Yang and J. H. H. Perk, Onsager’s star-triangle equation: Master key to integrability, In *Integrable system in quantum field theory and statistical mechanics*, Adv. Stud. Pure Math., 19, Kinokuniya Academic, Academic Press, Boston, MA (1989) 57–94.

- [4] H. Au-Yang and J.H.H. Perk, Eigenvectors the superintegrable model I: $\mathfrak{sl}_2$ generators, J. Phys. A: Math. Theor. 41 (2008) 275201; arXiv: 0710.5257; Eigenvectors in the superintegrable model II: ground state sector, arXiv: 0803.3029.
- [5] R. J. Baxter, Exactly solved models in statistical mechanics, Academic Press (1982).
- [6] R. J. Baxter, Chiral Potts model: eigenvalues of the transfer matrix, Phys. Lett. A 146 (1990) 110–114.
- [7] R. J. Baxter, Calculation of the eigenvalues of the transfer matrix of the chiral Potts model, *Proc. Fourth Asia-Pacific Physics Conference* (Seoul, Korea, 1990) Vol 1, World-Scientific, Singapore (1991) 42–58.
- [8] R. J. Baxter, Chiral Potts model with skewed boundary conditions, J. Stat. Phys. 73 (1993) 461–495.
- [9] R. J. Baxter, V.V. Bazhanov and J.H.H. Perk, Functional relations for transfer matrices of the chiral Potts model, Int. J. Mod. Phys. B 4 (1990) 803–870.
- [10] R. J. Baxter, J. H. H. Perk and H. Au-Yang, New solutions of the star-triangle relations for the chiral Potts model, Phys. Lett. A 128 (1988) 138–142.
- [11] V.V. Bazhanov and Yu.G. Stroganov, Chiral Potts model as a descendant of the six-vertex model, J. Stat. Phys. 59 (1990) 799–817.
- [12] E. Date, M. Jimbo, K. Miki and T. Miwa, Cyclic representations of $U_q(\mathfrak{sl}(n+1, \mathbf{C}))$ at $q^N = 1$, Publ. RIMS, Kyoto Univ. 27 (1991) 347–366.
- [13] C. DeConcini and V. G. Kac, Representations of quantum groups at roots of unity, in *Operator Algebra, Unitary Representations, Enveloping Algebras, and Invariant Theory*, Paris (1989) *Progress in Mathematics* 92, Birkhäuser, Boston, Massachusetts (1990) 471– 506.
- [14] L. D. Faddeev, How algebraic Bethe Ansatz works for integrable models, eds. A. Connes, K. Gawedzki and J. Zinn-Justin, *Quantum symmetries/ Symmetries quantiques*, Proceedings of the Les Houches summer school, Session LXIV, Les Houches, France, August 1-September 8, 1995, North-Holland (1998), 149–219.
- [15] N.S. Han and A. Honecker: Low-Temperature Expansions and Correlation Functions of the $\mathbf{Z}_3$ -Chiral Potts Model, J.Phys. A27 (1994) 9-22; BONN-HE-93-13, hep-th/9304083.
- [16] A. N. Kirillov and N. Yu. Reshetikhin, Exact solution of the integrable XXZ Heisenberg model with arbitrary spin: I. The ground state and the excitation spectrum, J. Phys. A: Math. Gen. 20 (1987) 1565 – 1595.
- [17] P. P. Kulish, N. Yu. Reshetikhin and E. K. Sklyanin, Yang Baxter equation and representation theory, Lett. Math. Phys. 5 (1981) 393–403.

- [18] V. B. Matveev and A. O. Simnov, Some comments on the solvable chiral Potts model, *Lett. Math. Phys.* 19 (1990) 179–185.
- [19] B. M. McCoy, J. H. H. Perk, S. Tang and C. H. Sah, Commuting transfer matrices for the four-state self-dual chiral Potts model with a genus-three uniformizing Fermat curve, *Phys. Lett. A* 125 (1987) 9–14.
- [20] B. M. McCoy and S. S. Roan, Excitation spectrum and phase structure of the chiral Potts model. *Phys. Lett. A* 150 (1990) 347–354.
- [21] S. S. Roan, The Q-operator for root-of-unity symmetry in six vertex model, *J. Phys. A: Math. Gen.* 39 (2006) 12303-12325; cond-mat/0602375.
- [22] S. S. Roan, Fusion operators in the generalized $\tau^{(2)}$ -model and root-of-unity symmetry of the XXZ spin chain of higher spin, *J. Phys. A: Math. Theor.* 40 (2007) 1481-1511; cond-mat/0607258.
- [23] S. S. Roan, The transfer matrix of superintegrable chiral Potts model as the Q-operator of root-of-unity XXZ chain with cyclic representation of $U_q(sl_2)$, *J. Stat. Mech.* (2007) P09021; arXiv: 0705.2856.
- [24] S. S. Roan, On the equivalent theory of the generalized $\tau^{(2)}$ -model and the chiral Potts model with two alternating vertical rapidities, arXiv: 0710.2764.
- [25] S. S. Roan, Bethe equation of $\tau^{(2)}$ -model and eigenvalues of finite-size transfer matrix of chiral Potts model with alternating rapidities, *J. Stat. Mech.* (2008) P10001; arXiv:0805.1585.
- [26] S. S. Roan, On $\tau^{(2)}$ -model in chiral Potts model and cyclic representation of quantum group $U_q(sl_2)$, *J. Phys. A: Math. Theor.* 42 (2009) 072003; arXiv:0806.0216.
- [27] S. S. Roan, Duality and symmetry in chiral Potts model, *J. Stat. Mech.* (2009) P08012; arXiv:0905.1924.
- [28] S. S. Roan, Eigenvectors of an arbitrary Onsager sector in superintegrable $\tau^{(2)}$ -model and chiral Potts model, arXiv:1003.3621.